Linear Equations

POINTS TO REMEMBER

- 1. Linear Equation : An equation which involves only one variable with highest power, is called a linear equation.
- 2. Solving the Equation

To solve an equation means to find the value of the variable which satisfies the equation.

3. Rules for solving a Linear Equation :

The equality of a linear equation is not changed when

- (i) the same number is added or subtracted from both sides of the equation.
- (ii) both sides of the equation are multiplied or divided by the same non zero number.
 - 4. Transposition :

When any term of the equation is taken from one side to the other, this process is called transposition.

5. Cross multiplication

If
$$\frac{a}{b} = \frac{c}{d}$$
 then $a \times d = b \times c$

This process is called the cross multiplication.

6. Word problems can be solved by means of equation by representing the unknown quantity as x, y, z etc. and then solve the equation so formed by the condition or conditions given in the problem :

Q. 1. 5x - 16 = 19 - 2xSol. 5x - 16 = 19 - 2x $\Rightarrow 5x + 2x = 19 + 16$ (By Transposition) Q. $\Rightarrow 7x = 35$ $\therefore x = \frac{35}{7} = 5$ Ans. Q. 2. $3x - \frac{1}{2}x = 1\frac{1}{2}$ Sol. $3x - \frac{1}{2}x = 1\frac{1}{2}$ 74 $\Rightarrow \frac{6x - x}{2} = \frac{3}{2}$ $\Rightarrow 6x - x = 3 \Rightarrow 5x = 3$

 $\therefore \quad x = \frac{3}{5} \quad \text{Ans.}$ Q. 3. $3\frac{3}{4}x = 5x - 2\frac{1}{2}$ Sol. $3\frac{3}{4}x = 5x - 2\frac{1}{2}$ $\Rightarrow \frac{15}{4}x = 5x - \frac{5}{2}$ Multiplying by 4, the LCM of 4 and 2 $\frac{15}{4}x \times 4 = 5x \times 4 - \frac{5}{2} \times 4$

75

15x = 20x - 10 $\Rightarrow 15x - 20x = -10$ (By transposition) $\Rightarrow -5x = -10$ $\therefore x = \frac{-10}{5} = 2$ Ans. **Q.4.** $3x-5=\frac{2x}{2}+9$ **Sol.** $3x - 5 = \frac{2x}{2} + 9$ $\Rightarrow 3x \times 3 - 5 \times 3 = \frac{2x}{3} \times 3 + 9 \times 3$ (Multiplying by 3) \Rightarrow 9x - 15 = 2x + 27 \Rightarrow 9x - 2x = 27 + 15 (By transposition) \Rightarrow 7x = 42 $\therefore x = \frac{42}{7} = 6$ Ans. **Q.5.** (x-4)(x+4) = (x+4)(x-7) + 33**Sol.** (x-4)(x+4) = (x+4)(x-7) + 33 $\Rightarrow x^2 - 4x + 4x - 16$ $= x^2 - 7x + 4x - 28 + 33$ $\Rightarrow x^2 - 16 = x^2 - 3x + 5$ $\Rightarrow x^2 - x^2 + 3x = 5 + 16$ (By transposition)

Sol. $\frac{y-4}{5} + \frac{y+2}{2} = 10$ Multiplying by 10, the LCM of 5 and 2 $\frac{(y-4)}{5} \times 10 + \frac{(y+2)}{2} \times 10 = 10 \times 10$ $\Rightarrow 2(y-4) + 5(y+2) = 100$ $\Rightarrow 2y-8+5y+10=100$ $\Rightarrow 2y + 5y = 100 + 8 - 10$ (By transposition) $\Rightarrow 7y = 98$ $\Rightarrow y = \frac{9.8}{7} = 14$ $\therefore y = 14$ Ans. **Q. 8.** $(x-1) = \frac{3}{4}(x+1) - \frac{1}{2}$ Sol. $(x-1) = \frac{3}{4}(x+1) - \frac{1}{2}$ Multiplying by 4, the L.C.M. of 4 and 2 $4(x-1) = \frac{3}{4}(x+1) \times 4 - \frac{1}{2} \times 4$ \Rightarrow 4x - 4 = 3x + 3 - 2 \Rightarrow 4x - 3x = 3 - 2 + 4 (By transposition) x = 5 Ans. Q.9. $\frac{2}{2}(x-3) = 1 - \frac{5}{6}(3x-4)$

$$\Rightarrow 3x = 21$$

$$\therefore x = \frac{21}{3} = 7 \text{ Ans.}$$

Q. 6. $(x-2)(x+3) = (x^2 - 4)$
Sol. $(x-2)(x+3) = x^2 - 4$

$$\Rightarrow x^2 + 3x - 2x - 6 = x^2 - 4$$

$$\Rightarrow x^2 + x - 6 = x^2 - 4$$

$$\Rightarrow x^2 + x - x^2 = -4 + 6$$

(By transposition)

$$\Rightarrow x = 2 \text{ Ans.}$$

Q. 7. $\frac{(y-4)}{5} + \frac{(y+2)}{2} = 10$

Sol. $\frac{2}{3}(x-3) = 1 - \frac{3}{6}(3x-4)$ Multiplying by 6, the LCM of 3 and 6 $6 \times \frac{2}{3}(x-3) = 1 \times 6 - 6 \times \frac{5}{6}(3x-4)$ $\Rightarrow 4(x-3) = 6 - 5(3x-4)$ $\Rightarrow 4x - 12 = 6 - 15x + 20$ $\Rightarrow 4x + 15x = 6 + 20 + 12$ (By transposition) $\Rightarrow 19x = 38$ $\therefore x = \frac{38}{10} = 2$ Ans.

76

Q. 10. $\frac{(x+7)}{2} = 1 + \frac{(3x-2)}{5}$ Sol. $\frac{x+7}{2} = 1 + \frac{3x-2}{5}$ Multiplying by 15, the LCM of 3 and 5 $15 \times \frac{(x+7)}{2} = 1 \times 15 + 15 \times \frac{(3x-2)}{5}$ $\Rightarrow 5(x+7) = 15+3(3x-2)$ $\Rightarrow 5x + 35 = 15 + 9x - 6$ \Rightarrow 5x - 9x = 15 - 6 - 35 $\Rightarrow -4x = -26 \Rightarrow x = \frac{-26}{4}$ $\therefore \qquad x = \frac{13}{2} = 6\frac{1}{2}$ Ans. Q. 11. $\frac{4}{5}\left(x+\frac{5}{8}\right)-\frac{2}{3}\left(x-\frac{1}{4}\right)=1\frac{1}{9}$ Sol. $\frac{4}{5}\left(x+\frac{5}{8}\right)-\frac{2}{3}\left(x-\frac{1}{4}\right)=1\frac{1}{9}$ $\Rightarrow \frac{4}{5} \frac{(8x+5)}{8} - \frac{2}{3} \frac{(4x-1)}{4} = \frac{10}{9}$ $\frac{8x+5}{10} - \frac{4x-1}{6} = \frac{10}{9}$ Multiplying by 90, the LCM of 10, 6 and 9, $90 \times \frac{8x+5}{10} - 90 \times \frac{4x-1}{6} = \frac{10}{9} \times 90$

 $\Rightarrow 2(x-2) + 15x = 36 - (x-5)$ 2x - 4 + 15x = 36 - x + 52x + 15x + x = 36 + 5 + 418x = 45 $\therefore x = \frac{45}{18} = \frac{5}{2} = 2\frac{1}{2}$ Ans. Q. 13. $4\frac{1}{3} - \frac{(3x-4)}{5} = \frac{(x-7)}{3}$ Sol. $\frac{13}{3} - \frac{3x-4}{5} = \frac{x-7}{2}$ Multiplying by 15, the LCM of 3 and 5. $15 \times \frac{13}{3} - 15 \times \frac{3x - 4}{5} = 15 \times \frac{x - 7}{3}$ $\Rightarrow 65 - 3(3x - 4) = 5(x - 7)$ $\Rightarrow \quad 65 - 9x + 12 = 5x - 35$ $\Rightarrow -9x - 5x = -35 - 65 - 12$ (By transposition) $\Rightarrow -14x = -112$ $\therefore x = \frac{-112}{-14} = 8$ Ans. Q.14. $\frac{(y-2)}{4} + \frac{1}{2} = y - \frac{(2y-1)}{2}$ Sol. $\frac{(y-2)}{4} + \frac{1}{3} = y - \frac{(2y-1)}{3}$ Multiplying 12, the LCM of 4 and 3

$$\Rightarrow 9 (8x + 5) - 15 (4x - 1) = 100$$

$$\Rightarrow 72x + 45 - 60x + 15 = 100$$

$$\Rightarrow 72x + 45 - 60x + 15 = 100$$

$$\Rightarrow 72x - 60x = 100 - 45 - 15$$

$$\Rightarrow 2x = 40$$

$$\therefore x = \frac{40}{12} = \frac{10}{3} = 3\frac{1}{3} \text{ Ans.}$$

$$2x = 40$$

$$\therefore x = \frac{40}{12} = \frac{10}{3} = 3\frac{1}{3} \text{ Ans.}$$

$$Q. 12. \quad \frac{(x - 2)}{3} + \frac{5x}{2} = 6 - \frac{(x - 5)}{6}$$

Multiplying by 6, the LCM of 3, 2 and 6

$$6 \times \frac{x - 2}{3} + 6 \times \frac{5x}{2} = 6 \times 6 - 6 \times \frac{x - 5}{6}$$

Multiplying by 6, the LCM of 3, 2 and 6

$$6 \times \frac{x - 2}{3} + 6 \times \frac{5x}{2} = 6 \times 6 - 6 \times \frac{x - 5}{6}$$

Sol.
$$(7x - 1) - \left[x - \frac{1 - x}{2}\right] = 5x + \frac{1}{2}$$

77

$$\Rightarrow 7x - 1 - x + \frac{1 - x}{2} = 5x + \frac{1}{2}$$

Multiplying by 2,

$$14x - 2 - 2x + \frac{(1 - x)}{2} \times 2 = 10x + \frac{1}{2} \times 2$$

$$\Rightarrow 14x - 2 - 2x + \frac{1 - x}{2} = 10x + \frac{1}{2} \times 2$$

$$\Rightarrow 14x - 2 - 2x + \frac{1 - x}{2} = 10x + 1$$

$$\Rightarrow 14x - 2x - x - \frac{10x}{10x} = 1 + 2 - 1$$

(By transposition)

$$\Rightarrow 14x - 13x = 3 - 1$$

$$\therefore x = 2$$
 Ans.
16. $\frac{(5x - 7)}{4} - \frac{(2x - 5)}{3} = \frac{5x}{6}$
Multiplying by 12, the LCM of 4, 3 and

$$6$$

$$12 \times \frac{5x - 7}{4} - \frac{12 \times \frac{2x - 5}{3}}{3} = 12 \times \frac{5x}{6}$$

$$\Rightarrow 3 (5x - 7) - 4 (2x - 5) = 10x$$

$$\Rightarrow 15x - 21 - 8x + 20 = 10x$$

$$\Rightarrow 15x - 8x - 10x = 21 - 20$$

(By transposition)

$$\Rightarrow 15x - 18x = 1$$

$$\Rightarrow -3x = 1$$

$$\therefore x = -\frac{1}{2}$$
 Ans.

	7 _ 5
18.	$\frac{7}{(x-4)} = \frac{5}{(x+2)}$
Sol.	$\frac{7}{x-4} = \frac{5}{x+2}$
	By cross multiplication,
	7(x+2) = 5(x-4)
	$\Rightarrow 7x + 14 = 5x - 20$
	$\Rightarrow 7x - 5x = -20 - 14 \Rightarrow 2x = -34$
	$\therefore x = \frac{-34}{2} = -17$ Ans.
19.	$\frac{x+1}{x-2} = \frac{x-2}{x-3}$
Sal	$\frac{x+1}{x-2} = \frac{x-2}{x-3}$
501.	BUCK III I
	By cross multiplication,
	(x+1)(x-3) = (x-2)(x-2)
	$\Rightarrow x^2 - 3x + x - 3 = x^2 - 2x - 2x + 4$
	$\Rightarrow x^2 - 3x + x - x^2 + 2x + 2x = 4 + 3$
	(By transposition)
	$\Rightarrow 5x - 3x = 7 \Rightarrow 2x = 7$
***	:. $x = \frac{7}{2} = 3\frac{1}{2}$ Ans.
	(2x-3) $(3x-1)$
Q. 20.	$\frac{(2x-3)}{(2x-1)} = \frac{(3x-1)}{(3x+1)}$
C.I	$\frac{2x-3}{2x-1} = \frac{3x-1}{3x+1}$
501.	
	By cross multiplication,

Q. 17. $\frac{5-4x}{3-2x} = 1\frac{6}{7}$ Sol. $\frac{5-4x}{3-2x} = \frac{13}{7}$ By cross multiplication, 7 (5-4x) = 13 (3-2x) $\Rightarrow 35-28x = 39-26x$ $\Rightarrow -28x + 26x = 39-35$ (By transposition) $\Rightarrow -2x = 4$ $\therefore \qquad x = \frac{4}{-2} = -2$ Ans.

Э

By cross multiplication, (2x-3) (3x + 1) = (3x - 1) (2x - 1) $\Rightarrow 6x^{2} + 2x - 9x - 3 = 6x^{2} - 3x - 2x + 1$ $\Rightarrow 6x^{2} + 2x - 9x - 6x^{2} + 3x + 2x = 1 + 3$ (By transposition) $\Rightarrow 7x - 9x = 4$ $\Rightarrow -2x = 4 \Rightarrow x = \frac{4}{-2}$ $\therefore x = -2 \text{ Ans.}$ Q. 21. $2 - \frac{(3-x)}{(x-1)} = \frac{(3x+4)}{(x+1)}$ Sol. $2 - \frac{3-x}{x-1} = \frac{3x+4}{x+1}$

78

 $\Rightarrow \frac{2x-2-3+x}{x-1} = \frac{3x+4}{x+1}$ $\Rightarrow \quad \frac{3x-5}{x-1} = \frac{3x+4}{x+1}$ By cross multiplication, -(3x-5)(x+1) = (3x+4)(x-1) \Rightarrow $3x^2 + 3x - 5x - 5$ $=3x^{2}-3x+4x-4$ $\Rightarrow 3x^2 + 3x - 5x + 3x^2 + 3x - 4x$ = -4 + 5 $\Rightarrow 6x - 9x = 1$ $\Rightarrow -3x = 1$ $\therefore x = -\frac{1}{2}$ Ans. Q. 22. $\frac{4}{(x-3)} + \frac{2}{(x-2)} = \frac{6}{x}$ Sol. $\frac{4}{r-3} + \frac{2}{r-2} = \frac{6}{r}$ $\Rightarrow \frac{4x-8+2x-6}{(x-3)(x-2)} = \frac{6}{x}$ $\Rightarrow \frac{6x-14}{x^2-2x-3x+6} = \frac{6}{x}$ $\Rightarrow \frac{6x-14}{x^2-5x+6} = \frac{6}{x}$ By cross multiplication, $x (6x - 14) = 6 (x^2 - 5x + 6)$

Multiplying each by 12, the LCM of 4, 2 and 3 $\frac{5y-11}{4} \times 12 + \frac{3y-7}{2} \times 12$ $=\frac{4y-7}{2} \times 12 + (y-1) 12$ $\Rightarrow 3 (5 y - 11) + 6 (3 y - 7)$ = 4 (4 v - 7) + 12 (v - 1) \Rightarrow 15 y - 33 + 18 y - 42 = 16 y - 28 + 12 y - 12 $\Rightarrow 15 y + 18 y - 16 y - 12 y$ = -28 - 12 + 33 + 42 \Rightarrow 33 y - 28 y = 75 - 40 \Rightarrow 5 y = 35 $\Rightarrow y = \frac{35}{5} = 7$ \therefore y = 7 Ans. **Q. 24.** Solve : $\frac{(2-x)}{2} - \frac{(x-3)}{3} = 1 - x$. Find y, when $\frac{1}{r} + \frac{1}{v} = 2$. Sol. $\frac{2-x}{2} - \frac{x-3}{3} = 1-x$ Multiplying by 6, the LCM of 2 and 3 $6 \times \frac{2-x}{2} - 6 \times \frac{x-3}{2} = 6 - 6x$

$$\Rightarrow 6x^{2} - 14x = 6x^{2} - 30x + 36$$

$$\Rightarrow 6x^{2} - 14x - 6x^{2} + 30x = 36$$

$$\Rightarrow 16x = 36$$

$$\Rightarrow 16x = 36$$

$$\Rightarrow x = \frac{36}{16} = \frac{9}{4}$$

$$\therefore x = 2\frac{1}{4} \text{ Ans.}$$
Q. 23. Solve :
$$\frac{5y - 11}{4} + \frac{3y - 7}{2} = \frac{4y - 7}{3} + y - 1$$
Sol. $\frac{5y - 11}{4} + \frac{3y - 7}{2} = \frac{4y - 7}{3} + y - 1$

$$y = \frac{6}{13}$$
Hence $x = 6$, and $y = \frac{6}{13}$ Ans.

79

Q. 25. If x = (k + 1), find the value of k, when,

$$\frac{1}{2}(5x-3) - \frac{1}{3}(2+9k) = \frac{1}{4}.$$

Sol. x = k + 1(*i*)

and
$$\frac{1}{2}(5x-3) - \frac{1}{3}(2+9k) = \frac{1}{4}$$
...(*ii*)

Substitution the value of x in (ii)

$$\frac{1}{2}[5(k+1)-3] - \frac{1}{3}(2+9k) = \frac{1}{4}$$
$$\Rightarrow \quad \frac{5}{2}(k+1) - \frac{3}{2} - \frac{2+9k}{3} = \frac{1}{4}$$

Multiplying by 12, the LCM of 2, 3 and 4.

$$12 \times \frac{5}{2}(k+1) - 12 \times \frac{3}{2} - 12 \times \frac{2+9k}{3}$$
$$= 12 \times \frac{1}{2}$$

4

$$\Rightarrow 30 (k+1) - 6 \times 3 - 4 (2+9k) = 3$$

$$\Rightarrow 30k + 30 - 18 - 8 - 36k = 3$$

$$\Rightarrow 30k - 36k = 3 - 30 + 18 + 8$$

$$\Rightarrow 6k = 29 - 30 = -1$$

$$k = \frac{-1}{-6} = \frac{1}{6} \text{Ans.}$$

EXERCISE 7 (B)

- Q. 1. Find a number, twice of which decreased by 7 gives 65.
- Sol. Let the number = x

Now according to the condition,

$$\Rightarrow \frac{1}{5}x = 23 - 8 = 15$$

$$\therefore \quad x = 15 \times \frac{5}{1} = 75$$

Hence number = 75 Ans.

Q. 3. Find a number such that one-fifth of it is less than by one-fourth of it by 3.

Sol. Let number = xAccording to the condition,

$$\frac{x}{5} = \frac{x}{4} - 3$$

$$\Rightarrow \frac{x}{4} - \frac{x}{5} = 3$$

$$\Rightarrow \frac{5x - 4x}{20} = 3 \implies \frac{x}{20} = 3$$

$$\therefore x = 3 \times 20 = 60$$

Hence number = 60 Ans.

Q. 4. Twice a number increased by 10 is 14 less than thrice the number. Find the number.

Sol. Let number
$$= x$$

According to the cor

According to the condition,

2x + 10 = 3x - 14

 $3x - 2x = 10 + 14 \implies x = 24$

Hence number = 24 Ans.

- Q. 5. Find three consecutive natural numbers whose sum is 120.
- Sol. Let first natural number = xsecond natural number = x + 1

2x - 7 = 65 2x = 65 + 7 = 72 $x = \frac{72}{2} = 36$ Ans. Check : $36 \times 2 - 7 = 72 - 7 = 65$ which is given. Hence number = 36 Q. 2. One-fifth of a number increased by 8 is equal to 23. Find the number. Sol. Let number = x According to the condition, $\frac{1}{5}x + 8 = 23$ and third natural number = x + 2According to the condition, x + x + 1 + x + 2 = 120 $\Rightarrow \quad 3x + 3 = 120$ $\Rightarrow \quad 3x = 120 - 3 = 117$ $\therefore \qquad x = \frac{117}{3} = 39$ \therefore first natural number = 39 and other numbers are 39, 40, 41 Ans. Q. 6. The difference of the squares of two consecutive even natural numbers is 92. Find the numbers.

80

- Sol. Let first even number = x then second even number = x + 2According to the condition, $(x + 2)^2 - x^2 = 92$ $\Rightarrow x^2 + 4x + 4 - x^2 = 92$ $\Rightarrow 4x + 4 = 92$
 - $\Rightarrow 4x = 92 4 = 88$

$$x = \frac{88}{4} = 22$$

...

...

Hence First even number = 22and second even number = 22 + 2= 24 Ans.

- Q. 7. Find two consecutive positive odd integers whose sum is 156.
- Sol. Let first odd number = 2x + 1Then second odd number = 2x + 3
- : According to the condition,
 - 2x + 1 + 2x + 3 = 156
 - \Rightarrow 4x + 4 = 156
 - $\Rightarrow 4x = 156 4 = 152$

$$x = \frac{152}{4} = 38$$

:. First odd number = 2x + 1= $2 \times 38 + 1 = 76 + 1 = 77$ and second odd number = 77 + 2 = 79Hence odd numbers are = 77, 79 Ans. Hence numbers are 64, 66 Ans.

- Q. 9. The sum of two numbers is 58 and their difference is 12. Find the numbers.
- Sol. Sum of two numbers = 58 Let first number = xthen second number = 58 - x

According to the condition,

x - (58 - x) = 12 $\Rightarrow x - 58 + x = 12$ $\Rightarrow 2x = 12 + 58$ $\Rightarrow 2x = 70$ $\therefore x = \frac{70}{2} = 35$

:. First number = 35and second number = 58 - 35 = 23Hence numbers are 35 and 23 Ans.

- Q. 10. Divide 88 into two parts such that when the larger is divided by the smaller, the quotient is 3 and the remainder is 4.
 - Sol. Sum of two parts = 88 Let larger part = xand smaller part = 88 - xQuotient = 3 and remainder = 4 We know that Dividend = Divisor \times Quotient + Remainder
 - : Larger part = smaller part $\times 3 + 4$
- Q. 8. Find two consecutive positive even integers whose sum is 130. Sol. Let first even positive integer = 2xthen second even number = 2x + 2According to the condition, 2x + 2x + 2 = 130 $\Rightarrow 4x + 2 = 130$ $\Rightarrow 4x = 130 - 2 = 128$ $\therefore x = \frac{128}{4} = 32$ \therefore First even number = $2x = 2 \times 32 = 64$ and second number = 2x + 2 = 64 + 2 = 66

 $\Rightarrow x = (88 - x) \times 3 + 4$ $\Rightarrow x = 264 - 3x + 4$ $\Rightarrow x + 3x = 268 \Rightarrow 4x = 268$ $\therefore x = \frac{268}{4} = 67$ $\therefore \text{ Larger part} = 67$ and smaller part = 88 - 67 = 21 Ans. Q. 11. The ages of A and B are in the ratio 9 : 4 Seven years hence, the ratio of their age will be 5 : 3. Find their present ages. Sol. Ratio in the present ages of A and B = 9 : 4

and age of father = 6x + 10Let A's age = 9x $= 6 \times 5 + 10 = 30 + 10 = 40$ years then B's age = 4xHence their present ages are 40 years and 7 years hence, 15 years Ans. A's age will be = 9x + 7Q. 13. Meena is five times as old as her son and B's age will be = 4x + 7Ashish. In 8 years time, Meena will be According to the condition, three times as old as Ashish. Find their $\frac{9x+7}{4x+7} = \frac{5}{3}$ present ages. Sol. Let present age of her son Ashish 3(9x+7) = 5(4x+7)= x years 27x + 21 = 20x + 35then Meena's age = 5x years \Rightarrow 7x - 20x = 35 - 21After 8 years, 7x = 14Age of Ashish will be = (x + 8) years \Rightarrow and age of Meena = (5x + 8) years $x = \frac{14}{7} = 2$ According to the condition, \therefore A's present age = $9x = 9 \times 2$ 5x + 8 = 3(x + 8)= 18 years 5x + 8 = 3x + 24and B's age = $4x = 4 \times 2 = 8$ years Ans. $\Rightarrow 5x - 3x = 24 - 8$ 10 years ago, a man was six times as old Q. 12. $\Rightarrow 2x = 16 \Rightarrow x = \frac{16}{2} = 8$ as his daughter. After 10 years, he will be twice as old as his daughter. Find their : Ashish's present age = 8 years present ages. and Meena's age = $5x = 5 \times 8$ Sol. 10 years ago, = 40 years Ans. Let age of daughter = x years Q. 14. A number consists of two digits. The digit then age of father = 6x years Present age of daughter = (x + 10) years

at ten's place is twice the digit at units place. The number formed by reversing the digits is 27 less than the original number. Find the original number.

age of daughter = (x + 10 + 10)= (x + 20) years and age of father = (6x + 10 + 10)= (6x + 20) years According to the condition, 6x + 20 = 2 (x + 20) $\Rightarrow 6x + 20 = 2x + 40$ $\Rightarrow 6x - 2x = 40 - 20 \Rightarrow 4x = 20$ $\therefore x = \frac{20}{4} = 5$ \therefore Present age of daughter = (x + 10) years = 5 + 10 = 15 years

and age of father = (6x + 10) years

and 10 years after,

Sol. Let digit at unit's place = x then digit at ten's place = 2x \therefore number = $x + (10 \times 2x)$ = x + 20x = 21xBy reversing the digits, the unit's digit = 2xand ten's digit = x \therefore Number = $2x + 10 \times x = 2x + 10x$ = 12xAccording to the condition, 12x = 21x - 27

 $\Rightarrow 12x - 21x = -27 \Rightarrow -9x = -27$ $\therefore \qquad x = \frac{-27}{-9} = 3$ $\therefore \text{ Original number} = 21x = 21 \times 3$

= 63 Ans.

- Q. 15. The ten's digit of a two-digit number exceeds its unit's digit by 5. The number itself is equal to 8 times the sum of the digits. Find the number.
 - Sol. Let unit's digit of the number = xthen ten's digit = x + 5

:. Number =
$$x + 10 (x + 5)$$

= x + 10x + 50 = 11x + 50

According to the condition,

$$11x + 50 = 8 (x + x + 5) = 8 (2x + 5)$$

$$\Rightarrow 11x + 50 = 16x + 40$$

$$\Rightarrow 11x - 16x = 40 - 50$$

$$\Rightarrow -5x = -10$$

$$\therefore \qquad x = \frac{-10}{-5} = 2.$$

Hence number = $11x + 50 = 11 \times 2 + 50$ = 22 + 50 = 72 Ans.

Q. 16. The denominator of a fraction is 4 more than its numerator. If 1 is subtracted from both the numerator and the denominator,

the fraction becomes $\frac{1}{2}$. Find the original fraction.

- Q. 17. The height of a triangle is 3 cm more than its base. If the area of the triangle is 10⁴ cm², find the lengths of its base and height.
 - Sol. Let base of a triangle (b) = x cmthen height (h) = (x + 3) cmand area $(A) = 104 \text{ cm}^2$ According to the condition,

$$\frac{1}{2}b \times h = A$$

 $\Rightarrow \frac{1}{2}x \times (x+3) = 104$ $\Rightarrow x (x+3) = 208$ $\Rightarrow x^2 + 3x - 208 = 0$ $\Rightarrow x^2 + 16x - 13x - 208 = 0$ {:: 3 = 16 - 13, and $-208 = 16 \times (-13)$ } $\Rightarrow x (x+16) - 13 (x+16) = 0$ $\Rightarrow (x+16) (x-13) = 0$ Either x + 16 = 0, then x = -16 which is not possible as it is in negative or x - 13 = 0 then x = 13 \therefore Base = 13 cm

and height = 13 + 3 = 16 cm Ans.

- Q. 18. Last year the prices of two houses were in the ratio 16 : 23. This year, the price of the first house has risen by 25% and that of the second by Rs. 5200 and the ratio of their new prices is 9 : 11. Find
- Sol. Let numerator of the fraction = xthen its denominator = x + 4

 $\therefore \text{ Fraction} = \frac{x}{x+4}$ According to the condition, $\frac{x-1}{x+4-1} = \frac{1}{2} \implies \frac{x-1}{x+3} = \frac{1}{2}$

$$\Rightarrow 2x - 2 = x + 3$$

(By cross multiplication) $2x - x = 3 + 2 \implies x = 5$ \therefore Fraction $= \frac{x}{x+4} = \frac{5}{5+4} = \frac{5}{9}$ Ans. ratio of their new prices is 9 : 11. Find their last year's prices. Sol. Let the price of first house = 16x then price of second house = 23x This year, the price of first house $= \frac{16x (100 + 25\%)}{100}$ $= 16x \times \frac{125}{100} = 20x.$ and price of second house = 23x + 5200 According to the condition, $\frac{20x}{23x + 5200} = \frac{9}{11}$

By cross multiplication, $\Rightarrow 11 \times 20x = 9 (23x + 5200)$ $\Rightarrow 220x = 207x + 46800$ $\Rightarrow 220x - 207x = 46800$ $\Rightarrow 13x = 46800 \Rightarrow x = \frac{46800}{13}$ $\therefore x = 3600$ $\therefore \text{ Price of first house} = 16x = 16 \times 3600$ = Rs. 57600and price of second house = 23x $= 23 \times 3600$ = Rs. 82800 Ans.Q. 19. There are 100 multiple-choice questions

- Q. 19. There are 100 multiple-choice questions in an engineering enterance examination. A candidate was given 5 marks for every correct answer and penalised 2 marks for every wrong answer. Pankaj answered all the questions and scored 241 marks. How many questions did he answer correctly?
 - Sol. Let no. of questions answered correctly

Sol. Let the distance Tetween school

then no. of questions which were wrongly attempted = 100 - xAccording to the condition, $x \times 5 - (100 - x) \times 2 = 241$

$$\Rightarrow 5x - 200 + 2x = 241$$

 $\Rightarrow 7x = 241 + 200 = 441$ $x = \frac{441}{7} = 63$ $\therefore \text{ No. of questions, he answered}$ correctly = 63 Ans. **Sol.** During a week, working hours = 40 No. of hours as overtime = xthen no. of hours for normal time = 40 - xPayment is made = Rs. 20 for a normal hour and Rs. 30 for overtime hour. Total payment received by a worker = Rs. 880According to the condition, $(40 - x) \times 20 + x \times 30 = 880$ 20 (40 - x) + 30x = 880800 - 20x + 30x = 880 $\Rightarrow 10x = 880 - 800 = 80$ $\therefore \quad x = \frac{80}{10} = 8$: No. of hours for overtime = 8 hours and normal hours = 40 - 8= 32 hours. Ans.

- Q. 21. The perimeter of a rectangular park is 80 m. If the length of the park be decreased by 2 m and breadth increased by 2 m, the area will be increased by 36 m². Find the original length and breadth of the park.
 - Sol. Perimeter of a rectangular park = 80 m

:. Length + breadth = $\frac{80}{2}$ = 40 m. Let length of park = x m then breadth = (40 - x) m

Q. 20. A worker in a factory is paid Rs. 20 per hour for normal work and Rs. 30 per hour for overtime work. During a week, he worked for 40 hours, out of which x hours was overtime. If he receives Rs. 880 in all, write an equation in x. Solve it and hence find the number of hours of his normal work during the week. and area = $lb = x (40 - x) m^2$ According to the condition, (x - 2) [40 - x + 2] = x (40 - x) + 36 $(x - 2) (42 - x) = 40x - x^2 + 36$ $42x - x^2 - 84 + 2x = 40x - x^2 + 36$ $\Rightarrow 42x - x^2 + 2x - 40x + x^2 = 36 + 84$ $\Rightarrow 44x - 40x = 120 \Rightarrow 4x = 120$ $\therefore x = \frac{120}{4} = 30$ \therefore Length of park = 30 m and breadth = 40 - 30 = 10 m Ans.

84

Q. 22. A man invested Rs. 5000, a part of it at 12% p.a and the rest at 14% p.a. If he received a total interest of Rs. 636, how much did he invest at 14% p.a.? Sol. Total sum invested = Rs. 5000 Let investment at 14% = Rs. x then investment at 12% = Rs. (5000 - x) Total interest received = Rs. 636 According to the condition, $\frac{x \times 14 \times 1}{100} + \frac{(5000 - x) \times 12 \times 1}{100} = 636$ $\Rightarrow \frac{14x}{100} + \frac{12(5000 - x)}{100} = 636$ $\Rightarrow 14x + 12(5000 - x) = 63600$

 $\Rightarrow 14x + 60000 - 12x = 63600$ $\Rightarrow 14x - 12x = 63600 - 60000$

$$2x = 3600$$

a stream bark n

$$x = \frac{3600}{2} = 1800$$

0. 21. T

Hence sum invested on 14% p.a.

= Rs. 1800 Ans.

- Q. 23. By selling a T.V. set for Rs. 27600, a trader makes a profit of 15%. What is the cost price of the set ?
 - Ans. Let CP of the T.V. set = Rs. xGain = 15%

Sol. Total cost of desk and chair = Rs. 477 Let cost of chair = x

then cost of desk = $\frac{x \times (100 + 12)}{100}$

$$=\frac{112}{100}x$$

According to the condition,

$$x + \frac{112}{100}x = 477$$

$$\Rightarrow 100x + 112x = 47700$$

$$\Rightarrow 212x = 47700$$

$$\Rightarrow x = \frac{47700}{212} = 225$$

$$\therefore \text{ Cost of chair = Rs. 225}$$

and cost of desk = Rs. 477 - 222

$$= \text{Rs. 252 Anse}$$

- Q. 25. Amit walks from his house to school at a speed of 3 kmph and returns back at a speed of 4 kmph. If he takes 42 minutes for the whole journey, find the distance between his house and the school.
 - Sol. Let the distance between school and house = x km

Total time taken = 42 minutes

$$=\frac{42}{60}$$
 hours

25

According to the condition,

S.P. = Rs. 27600 According to the condition, $SP = \frac{CP(100 + gain\%)}{100}$ $27600 = \frac{x \times (100 + 15)}{100} = \frac{115x}{100}$ $\therefore 115x = 27600 \times 100 = 2760000$ $x = \frac{2760000}{115} = \text{Rs. 24000}$ $\therefore \text{ Cost price of the set} = \text{Rs. 24000 Ans.}$ Q. 24. The total cost of a desk and a chair is Rs. 477. If the desk costs 12% more than the chair, find the cost of each. $\frac{x}{3} + \frac{x}{4} = \frac{42}{60} \left[\because \text{Time} = \frac{\text{Distance}}{\text{Speed}} \right]$ $\Rightarrow \frac{4x + 3x}{12} = \frac{42}{60}$ $\Rightarrow \frac{7x}{12'} = \frac{42}{60}$ $\Rightarrow \frac{7x}{12'} = \frac{42}{60}$ $\Rightarrow x = \frac{42}{60} \times \frac{12}{7} = \frac{6}{5}$ $\therefore \text{Distance between his house and school}$

 $=\frac{6}{5}\,\mathrm{km}=1.2\,\mathrm{km}\,\mathrm{Ans.}$



Q. 26. A man covers a distance of 184 km in 3 hours 30 minutes, partly by bus and partly by car. If their speeds be 48 kmph and 60 kmph respectively, find the distance covered by bus.

Sol. Total distance = 184 km.

Total time taken = 3 hour 30 minutes

$$=3\frac{1}{2}=\frac{7}{2}$$
 hours.

Let the distance covered by bus = x kmthen the distance covered by car

=(184 - x) km

According to the condition,

$$\frac{x}{48} + \frac{184 - x}{60} = \frac{7}{2}$$
$$\left[\because \text{Time} = \frac{\text{Distance}}{\text{Speed}}\right]$$

Multiplying by 240, the LCM of 48, 60 and 2

 $240 \times \frac{x}{48} + 240 \times \frac{184 - x}{60} = \frac{7}{2} \times 240$ $\Rightarrow 5x + 4 (184 - x) = 840$ \Rightarrow 5x + 736 - 4x = 840 $\Rightarrow 5x - 4x = 840 - 736$ x = 104 \therefore Distance covered by bus = 104 km Ans.

4x + 12 = 5x - 154x - 5x = -15 - 12-x = -27x = 27

Hence speed of steamer in still water = 27 kmph Ans.

- Q. 28. How much pure alcohol be added to a 400 ml, 15% solution to make it contain 32% alcohol?
 - Sol. Solution = 400 ml. and alcohol = 15%
 - \therefore Alcohol in the solution = $\frac{400 \times 25}{100}$

= 60 ml.

Let x ml of alcohol be added to make the solution 32%

According to the condition,

 $\left(\frac{60+x}{400+x}\right) \times 100 = 32$ $\Rightarrow \frac{60+x}{400+x} = \frac{32}{100}$ 100 (60 + x) = 32 (400 + x)(By cross multiplication) 6000 + 100x = 12800 + 32x100x - 32x = 12800 - 600068x = 6800

Q. 27. A steamer goes down stream and covers the distance between two ports in 4 hours white it covers the same distance upstream in 5 hours. If the speed of the stream be 3 kmph, find the speed of the steamer in still water.

Sol. Let the speed of steamer in still water = x kmphSpeed of stream = 3 kmph \therefore Speed of down stream = (x + 3) kmph and speed of upstream = (x - 3) kmph According to the condition, $(x+3) \times 4 = (x-3) \times 5$

 $x = \frac{6800}{68} = 100$: 100 ml of alcohol is to be added Ans. Q. 29. In a class room, there are x seats. If each student in the class occupies one seat, then 9 students remain standing and if 2 students occupy one seat, then 7 seats are left un-occupied. Find the number of seats in the class room and the number of students in the class.

Sol. No. of seats = x

If one student occupies one seat then number of students = x + 9

If two students occupy one seat then

86

number of students = 2(x - 7)

- $\therefore x+9=2(x-7)$
- $\Rightarrow 2x 14 = x + 9$
- $\Rightarrow 2x x = 14 + 9$
- \Rightarrow x = 23

 \therefore No. of seats = 23

and no. of students = x + 9 = 23 + 9

= 32 Ans.

Q. 30. In a shooting competition, a marks man receives Rs. 2 if he hits the mark and pays Re. 1 if he misses it. He tried 60 shots and was paid Rs. 18. How many times did he hit the mark?

Sol. No. of shots = 60 and amount received

= Rs. 18

Let no. of shots hit by him = xthen no. of shots missed = 60 - xAccording to the condition, $x \times 2 - (60 - x) \times 1 = 18$ 2x - 60 + x = 18 \Rightarrow 3x = 18 + 60 $\Rightarrow 3x = 78 \Rightarrow x = \frac{78}{3} = 26$ \therefore No. of shots he hit = 26 Ans.

EXERCISE 7 (C)

From (ii), substitute the value of x in (i) $y+3+y-7=0 \implies 2y-4=0$ Substituting the value of y in (i), $x+4x-5=0 \implies 5x-5=0$ $\Rightarrow 5x = 5 \Rightarrow x = \frac{5}{5} = 1$ Now substituting the value of x in (*iii*), $y = 4 \times 1 = 4$ Hence x = 1, y = 4 Ans. **Q. 2.** x + y - 5 = 0, y - 2x = 2x. **Sol.** x + y - 5 = 0...(i) y-2x=2x...(*ii*) From (ii) y = 2x + 2x = 4x...(iii) $x = 1 - 2 \times (-2) = 1 + 4 = 5$ Hence x = 5, y = -2 Ans. **O.3.** x + 2y - 1 = 0, 3x - y - 17 = 0**Sol.** x + 2y - 1 = 0...(i) 3x - y - 17 = 0...(*ii*) From (i) x = 1 - 2y ...(iii) Substituting the value of x in (*ii*), 3(1-2y)-y-17=0 \Rightarrow 3-6y-y-17=0 $\Rightarrow -7y - 14 = 0 \Rightarrow 7y + 14 = 0$ \Rightarrow 7 y = -14 \Rightarrow y = $\frac{-14}{7}$ = -2

- Q.1. Solve the following simultaneous equations :
- x + y 7 = 0, y + 3 = x.Sol. x + y - 7 = 0 ...(i) y + 3 = x ...(ii)

If two students eccapy analeast then

A steamer goes down stream and covers Substituting the value of y in (iii) $x = 1 - 2 \times -2$ $\Rightarrow x = 1 + 4$ $\Rightarrow x = 5$ Hence x = 5, y = -2 Ans.

s Speed of stream = 3 kmph.

ident (6 - 4) + incenter of Long bus - (t- 3) kmph

87

Q. 4.	5x + 4y - 4 = 0, x - 20 = 12y
Sol.	5x + 4y - 4 = 0(<i>i</i>)
	x - 20 = 12 y(<i>ii</i>)
	From (<i>ii</i>), $x = 12 y + 20$ (<i>iii</i>)
	Substituting the value of x in (i)
	5(12y+20) + 4y - 4 = 0
	$\Rightarrow 60y + 100 + 4y - 4 = 0$
	$\Rightarrow 64y + 96 = 0 \Rightarrow 64y = -96$
	-96 3
	$\Rightarrow \qquad y = \frac{-96}{64} = -\frac{3}{2}$
	Substituting the value of y in (<i>iii</i>)
	Substituting the value of vin (a
	$x = 12 \times \left(\frac{-3}{2}\right) + 20$
	= -18 + 20 = 2
	Hence x P 2, 4 Pus.
	Hence $x = 2$, $y = \frac{-3}{2}$ Ans.
0.5.	x + 2y + 9 = 0, 3x + 4y + 17 = 0
	x + 2y + 9 = 0(<i>i</i>)
	3x + 4y + 17 = 0(<i>ii</i>)
	From (i) $x = -2y - 9$ (iii)
	Substituting the value of x in (ii)
	3(-2y-9) + 4y + 17 = 0
	$\Rightarrow -6y - 27 + 4y + 17 = 0$

Substituting the value of y in (i)

$$2x + 3\left(\frac{5x - 20}{8}\right) = 23$$

$$2x + \frac{15x - 60}{8} = 23$$

$$16x + 15x - 60 = 184$$
(Multiplying by 8)

$$31x = 184 + 60 = 244$$

$$x = \frac{244}{31} = 7\frac{27}{31}$$
Substituting the value of x in (*iii*)


$$\left[5 \times \frac{244}{31} - 20\right] = \frac{1220}{31} - 20$$

$$= \frac{1220 - 620}{31 \times 8} = \frac{600}{31 \times 8} = \frac{75}{31}$$
$$= 2\frac{13}{31}$$

8

Hence $x = 7\frac{27}{31}$ and $y = 2\frac{13}{31}$ Ans. **Q.** 7. 3x + 7y = 15, 5x - 129 = 23y**Sol.** 3x + 7y = 15...(i) ...(ii) 5x - 129 = 23yFrom (*ii*) 23 y = 5 x - 129

$$y = \frac{5x - 129}{...(iii)}$$



 \Rightarrow 23 Substituting the value of y in (i) $3x + 7\left(\frac{5x - 129}{23}\right) = 15$ $\Rightarrow \quad 3x + \frac{35x - 903}{23} = 15$ 69x + 35x - 903 = 345= (Multiplying by 23) \Rightarrow 104 x = 345 + 903 \Rightarrow 104 x = 1248 $\Rightarrow \qquad x = \frac{1248}{104} = 12$

88

Substituting the value of x in (iii), $y = \frac{5 \times 12 - 129}{23} = \frac{60 - 129}{23} = \frac{-69}{23}$ = - 3 Hence x = 12 and y = -3 Ans. **Q.8.** 3 - (x - 5) = y + 2, 2(x + y) = 4 - 3ySol. 3 - (x - 5) = y + 23 - x + 5 = y + 2 $8 - x = y + 2 \implies x + y = 8 - 2$...(i) $\Rightarrow x + y = 6$

$$2 (x + y) = 4 - 3 y$$

$$\Rightarrow 2x + 2y = 4 - 3 y$$

$$\Rightarrow 2x + 2y + 3y = 4$$

$$\Rightarrow 2x + 5y = 4$$

$$\therefore 2x + 5y = 4$$

$$\therefore 2x + 5y = 4$$

$$\therefore (ii)$$

From (1) $x = 6 - y$

$$\dots (iii)$$

Substituting the value of x in (*ii*),

$$2 (6 - y) + 5 y = 4$$

$$12 - 2 y + 5 y = 4$$

$$3 y = 4 - 12 = -8$$

$$y = \frac{-8}{3}$$

Substituting the value of y in (*iii*),

$$x = 6 - \left(-\frac{8}{3}\right) = 6 + \frac{8}{3}$$

From (ii),

$$x = \frac{2y+7}{5} \qquad \dots (iii)$$

Substituting the value of x in (i),

$$8\left(\frac{2y+7}{5}\right) - 3y = 12$$

$$\Rightarrow \frac{16y+56}{5} - 3y = 12$$

$$\Rightarrow 16y+56 - 15y = 60$$
(Multiplying by 5)
$$\Rightarrow 16y - 15y = 60 - 56$$

$$y = 4$$
Substituting the value of y in (*iii*),
$$x = \frac{2 \times 4 + 7}{5} = \frac{8 + 7}{5} = \frac{15}{5} = 3$$
Hence $x = 3, y = 4$ Ans.

Q. 10.
$$x - \frac{2}{3}y = \frac{8}{3}, \frac{2x}{5} - y = \frac{7}{5}$$

Sol. $x - \frac{2}{3}y = \frac{8}{3}$...(*i*)
 $\frac{2x}{5} - y = \frac{7}{5}$...(*ii*)
From (*i*)
 $x = \frac{8}{3} + \frac{2}{3}y = \frac{8 + 2y}{3}$...(*iii*)

$$= \frac{18+8}{3} = \frac{26}{3}$$

Hence $x = \frac{26}{3}, y = \frac{-8}{3}$ Ans.
Q. 9. $2x - \frac{3y}{4} = 3, 5x = 2y + 7$
Sol. $2x - \frac{3y}{4} = 3$
 $\Rightarrow 8x - 3y = 12$
(Multiplying by 4)
 $8x - 3y = 12$
 $5x = 2y + 7$...(i)

Substituting the value of y in (*ii*),

3

(iii)

 $\frac{2}{5}\left(\frac{8+2y}{3}\right) - y = \frac{7}{5}$ $\Rightarrow \quad \frac{16+4y}{15} - y = \frac{7}{5}$ \Rightarrow 16+4y-15y=21 (Multiplying by 15, the L.C.M. of 15, 5) $\Rightarrow -11 y = 21 - 16 = 5$ $y = \frac{-5}{11}$

Substituting the value of y in (iii),

89

$$x = \frac{8+2\left(\frac{-5}{11}\right)}{3} = \frac{8-\frac{10}{11}}{3}$$

$$=\frac{88-10}{11\times3}=\frac{78}{11\times3}$$

$$x = \frac{26}{11}$$

Hence
$$x = \frac{26}{11}$$
, $y = \frac{-5}{11}$ Ans.

Q. 11.
$$\frac{3}{5}x - \frac{2}{3}y + 1 = 0, \frac{2}{5}x + \frac{1}{3}y - 4 = 0$$

Sol.
$$\frac{3}{5}x - \frac{2}{3}y + 1 = 0$$

 $\Rightarrow 15 \times \frac{3}{5}x - 15 \times \frac{2}{2}y + 15 = 0$ (Multiplying by 15, the L.C.M. of 5, 3) $\Rightarrow 9x - 10y + 15 = 0$...(i)

Again
$$\frac{2}{5}x + \frac{1}{3}y - 4 = 0$$

 $15 \times \frac{2}{5}x + 15 \times \frac{1}{3}y - 4 \times 15 = 0$ (Multiplying by 15, the L.C.M. of 5, 3) 6x + 5y - 60 = 0....(*ii*) Multiply (i) by 2 and (ii) by 3, 18 x - 20 y + 30 = 0

Q. 12. $\frac{x}{7} + \frac{y}{3} = 5, \frac{x}{2} - \frac{y}{9} = 6$ **Sol.** $\frac{x}{7} + \frac{y}{3} = 5$ Multiplying by 21, the L.C.M. of 7, 3 $21 \times \frac{x}{7} + 21 \times \frac{y}{3} = 21 \times 5$ 3x + 7y = 105 ...(*i*) Again $\frac{x}{2} - \frac{y}{9} = 6$ Multiplying by 18, the L.C.M. of 2 and 9, $18 \times \frac{x}{2} - 18 \times \frac{y}{9} = 6 \times 18$ 9x - 2y = 108...(ii) Multiplying (i) by 3 and (ii) by 1 9x + 21y = 3159x - 2y = 108+ 120- CSI -Subtracting, we get, 23 y = 207

$$y = \frac{207}{23} = 9$$

Substituting the value of y in (i), $3x + 7 \times 9 = 105 \implies 3x + 63 = 105$ 3x = 105 - 63 = 42 $x = \frac{42}{3} = 14$ Hence x = 14, y = 9 Ans. **Q. 13.** $\frac{x}{3} + \frac{y}{4} = 11$, $\frac{5x}{6} - \frac{y}{2} + 7 = 0$ **Sol.** $\frac{x}{3} + \frac{y}{4} = 11$ Multiplying by 12, the L.C.M. of 3 and $12 \times \frac{x}{3} + 12 \times \frac{y}{4} = 12 \times 11$ 4x + 3y = 132...(i)

18 x + 15 y - 180 = 0

lence x = + and =

Subtracting, we get, $-35y + 210 = 0 \implies -35y = -210$ $y = \frac{-210}{35} = 6$ Substituting the value of y in (i), 9x - 10(6) + 15 = 09x - 60 + 15 = 0 \Rightarrow $\Rightarrow 9x - 45 = 0 \Rightarrow 9x = 45$ $\therefore \quad x = \frac{45}{9} = 5$ Hence x = 5, y = 6 Ans. Downloaded from https:// www.studiestoday.com

90

Again $\frac{5x}{6} - \frac{y}{2} + 7 = 0$ Multiplying by 6, the L.C.M. of 6 and 3 $6 \times \frac{5}{6}x - 6 \times \frac{y}{3} + 6 \times 7 = 0$ 5x - 2v + 42 = 05x - 2y = -42...(*ii*) Multiply (i) by 5 and (ii) by 4, 20x + 15y = 66020x - 8y = -168Vicitiphy +g by 18+the L_C.M. of 2. and Subtracting, we get, 23 y = 828 $y = \frac{828}{22} = 36$ Substituting the value of y in (i), $4x + 3 \times 36 = 132$ \Rightarrow 4x + 108 = 132 \Rightarrow 4 x = 132 - 108 = 24 $\Rightarrow x = \frac{24}{4} = 6$ Hence x = 6, y = 36 Ans. Q. 14. $\frac{x}{6} + 6 = y$, $\frac{3x}{4} = 1 + y$ **Sol.** $\frac{x}{6} + 6 = y \implies y = \frac{x + 36}{6}$ $\frac{3x}{4} = 1 + y \implies y = \frac{3x}{4} - \dots(ii)$ From (i) and (ii) $\frac{x+36}{6} = \frac{3x}{4} - 1$ $\frac{x+36}{6} = \frac{3x-4}{4}$ By cross multiplication, 4(x+36) = 6(3x-4)4x + 144 = 18x - 244x - 18x = -24 - 144

-14x = -168 $x = \frac{-168}{14} = 12$ Substituting the value of x in (i), $y = \frac{12 + 36}{6} = \frac{48}{6} = 8$ x = 12, y = 8 Ans. Hence Q. 15. $x - y = \frac{9}{10}, \frac{11}{2(x + y)} = 1$ Sol. $x - y = \frac{9}{10}$...(i) $\frac{11}{2(x+y)} = 1 \implies 2(x+y) = 11$ $x + y = \frac{11}{2}$...(ii) Adding (i) and (ii), $2x = \frac{9}{10} + \frac{11}{2} = \frac{9+55}{10} = \frac{64}{10}$ $x = \frac{64}{10 \times 2} = \frac{16}{5}$ Subtracting (i) from (ii), $2y = \frac{11}{2} - \frac{9}{10} = \frac{55 - 9}{10} = \frac{46}{10}$ $y = \frac{46}{10 \times 2} = \frac{23}{10}$

Hence $x = \frac{16}{5}$ and $y = \frac{23}{10}$ Ans. Q. 16. $\frac{x}{2} + y = 0.8$, $\frac{7}{x + \frac{y}{2}} = 10$ Sol. $\frac{x}{2} + y = 0.8$ $\Rightarrow x + 2y = 1.6$...(*i*) And $\frac{7}{x + \frac{y}{2}} = 10$ $\Rightarrow x + \frac{y}{2} = \frac{7}{10} = 0.7$...(*ii*)

Subtracting (ii) from (i)

$$2y - \frac{y}{2} = 1 \cdot 6 - 0 \cdot 7 \implies \frac{3}{2}y = 0 \cdot 9$$
$$y = \frac{0 \cdot 9 \times 2}{3} = 0 \cdot 6 = \frac{3}{5}$$

Substituting the value of y in (i)

$$x + 2 \times \frac{3}{5} = \frac{16}{10} \implies x + \frac{6}{5} = \frac{8}{5}$$
$$x = \frac{8}{5} - \frac{6}{5} = \frac{2}{5}$$
Hence $x = \frac{2}{5}$ and $y = \frac{3}{5}$ Ans

Q. 17. Solve the following simultaneous equations :

5

$$4x + \frac{x - y}{8} = 17, \ x + 2y = \frac{y - 2}{3} - 2$$

5

Sol.
$$4x + \frac{x - y}{8} = 17$$

 $\Rightarrow 32 x + x - y = 136$
 $\Rightarrow 33 x - y = 136$...(i)
 $x + 2y = \frac{y - 2}{3} - 2$

$$\Rightarrow 3x + 6y = y - 2 - 6$$

$$\Rightarrow 3x + 6y - y = -8$$

$$\Rightarrow 3x + 5y = -8 \dots (ii)$$

From (i)

Q. 18. $\frac{7+x}{5} - \frac{2x-y}{4} = 3y - 5$ $\frac{4x-3}{6} + \frac{5y-7}{2} = 18 - 5x$ Sol. $\frac{7+x}{5} - \frac{2x-y}{4} = 3y - 5$ 4(7+x) - 5(2x-y) = 20(3y-5)(Multiplying by 20, the L.C.M. of 5 and 4) 28 + 4x - 10x + 5y = 60y - 100 $\Rightarrow 4x - 10x + 5y - 60y = -100 - 28$ $\Rightarrow -6x - 55y = -128$ $\Rightarrow 6x + 55y = 128$...(i) and $\frac{4x-3}{6} + \frac{5y-7}{2} = 18 - 5x$ $6 \times \frac{4x-3}{6} + 6 \times \frac{5y-7}{2} = 6(18-5x)$ (Multiplying by 6, the L.C.M. of 6 and 2) (4x-3)+3(5y-7)=108-30x4x - 3 + 15y - 21 = 108 - 30x4x + 15y + 30x = 108 + 21 + 334 x + 15 y = 132 ...(*ii*) Multiplying (i) by 3 and (ii) by 11 18x + 165y = 384374 x + 165 y = 1452

> Subtracting, we get, -356 x = -1068

y = 33x - 136...(*iii*) Substituting the value of y in (*ii*), 3 x + 5 (33 x - 136) = -8 ⇒ 3 x + 165 x - 680 = -8 ⇒ 168 x = -8 + 680 = 672 $x = \frac{672}{168} = 4$ Substituting the value of x in (*iii*), y = 33 x - 136 = 33 × 4 - 136 = 132 - 136 = -4 ∴ x = 4, y = -4 Ans. $x = \frac{-1068}{-356} = 3$ Substituting the value of x in (i), $6 \times 3 + 55 \ y = 128$ $\Rightarrow 18 + 55 \ y = 128$ $\Rightarrow 55 \ y = 128 - 18 = 110$ $\therefore \qquad y = \frac{110}{55} = 2$ Hence $x = 3, \ y = 2$ Ans. Q. 19. 6x + 5y = 7x + 3y + 1 = 2(x + 6y - 1)Sol. 6x + 5y = 7x + 3y + 1 $\Rightarrow 6x + 5y - 7x - 3y = 1$

92

-x+2y=1x - 2y = -1...(i) 7x + 3y + 1 = 2(x + 6y - 1) \Rightarrow 7x+3y+1=2x+12y-2 7x + 3y - 2x - 12y = -2 - 15x - 9y = -3...(*ii*) From (i) x = 2 y - 1...(*iii*) Substituting the value of x in (*ii*), 5(2y-1)-9y=-3 \Rightarrow 10 y - 5 - 9 y = -3 10 y - 9 y = -3 + 5v = 2Substituting the value of y in (iii), $x = 2 \times 2 - 1 = 4 - 1 = 3$ Hence x = 3, y = 2 Ans. **Q. 20.** 103 x + 51 y = 617, 97 x + 49 y = 583**Sol.** 103 x + 51 y = 617...(i) 97 x + 49 y = 583...(*ii*) Multiplying (i) by 49 and (ii) by 51, 5047 x + 2499 y = 302334947 x + 2499 y = 29733

Subtracting, 100 x = 500 $x = \frac{500}{100} = 5$ Adding, $17x = 51 \implies x = \frac{51}{17} = 3$ Substituting the value of x in (i), Substituting the value of x in (i), $103 \times 5 + 51 y = 617$ 515 + 51 y = 617 $4 \times 3 + \frac{6}{3} = 15$ $\Rightarrow 51 y = 617 - 515$ $12 + \frac{6}{v} = 15$ $y \Rightarrow 51 y = 102 \Rightarrow y = \frac{102}{51} = 2$ $\Rightarrow \frac{6}{-} = 15 - 12 = \frac{3}{-12}$ Hence x = 5, y = 2 Ans. **Q. 21.** 23 x - 29 y = 98, 29 x - 23 y = 110(By cross multiplicat **Sol.** 23 x - 29 y = 98 ...(*i*) $\therefore y = \frac{6}{2} = 2$ 29 x - 23 y = 110Adding (i) and (ii), we get Hence x = 3, y = 2 Ans.

52 x - 52 y = 208 $52 (x - y) = 208 \implies x - y = \frac{208}{52}$ x - y = 4Subtracting (ii) from (i) $-6x - 6y = -12 \implies 6x + 6y = 12$ $6 (x + y) = 12 \implies x + y = \frac{12}{6}$ x + y = 2Adding (iii) and (iv), we get, $2x=6 \implies x=3$ Subtracting (iii) from (iv) $2y = -2 \implies y = -1$ • Hence x = 3, y = -1 Ans. Q. 22. $4x + \frac{6}{v} = 15, 3x - \frac{4}{v} = 7$ **Sol.** $4x + \frac{6}{v} = 15$ $3x - \frac{4}{3} = 7$ Multiplying (i) by 2 and (ii) by 3,

$$8x + \frac{12}{y} = 30$$

9x - $\frac{12}{y} = 21$

93

		-
Q. 23.	$\frac{2}{x} + \frac{2}{3y} = \frac{1}{6}, \frac{3}{x} + \frac{2}{y} = 0$	
Sol.	$\frac{2}{x} + \frac{2}{3y} = \frac{1}{6}$.(i)
•	$\frac{3}{x} + \frac{2}{y} = 0$	
	Multiplying (i) by 3 and (ii) by 2,	
	$\frac{6}{x} + \frac{6}{3y} = \frac{3}{6}$	
	$\Rightarrow \qquad \frac{6}{x} + \frac{2}{y} = \frac{1}{2}$	
	- x y 2	
	$\frac{6}{-} + \frac{4}{-} = 0$	
	x y	
n (2),	Subtracting, $-\frac{2}{v} = \frac{1}{2}$	
	$\Rightarrow y = -4$	
	Substituting the value of y in (i) ,	
	$\frac{2}{2} + \frac{2}{2} = \frac{1}{2}$	
	$x^{+}3(-4)^{-}6$	
	$\frac{2}{2} + \frac{2}{2} = \frac{1}{2}$	

-12

 $\frac{2}{x} - \frac{1}{6} = \frac{1}{6}$


 $\frac{15}{2x} + \frac{10}{3y} = 25$ $\frac{15}{2x} - \frac{9}{2y} = \frac{3}{2}$

Subtracting, we get,

 $\frac{10}{3y} + \frac{9}{2y} = 25 - \frac{3}{2} = \frac{47}{2}$ $\frac{20}{6v} + \frac{27}{6v} = \frac{47}{2} \implies \frac{47}{6v} = \frac{47}{2}$ $\Rightarrow 6y \times 47 = 47 \times 2$ $y = \frac{47 \times 2}{6 \times 47} = \frac{1}{3}$ ⇒ Substituting the value of y in (i), $\frac{3}{2x} + \frac{3}{3 \times \frac{1}{2}} = 5 \implies \frac{3}{2x} + 2 = 5$ $\frac{3}{2x} = 5 - 2 = 3 \implies 6x = 3$ $x = \frac{3}{6} = \frac{1}{2}$ Hence $x = \frac{1}{2}, y = \frac{1}{3}$ Ans. Q. 25. $5x - 9 = \frac{1}{v}, x + \frac{1}{v} = 3$

...(i)

...(ii)



94

 $5 \times 2 - 9 = \frac{1}{y} \implies \frac{1}{y} = 10 - 9 = 1$ $\Rightarrow y=1$ Hence x = 2, y = 1 Ans. **Q. 26.** x + y = 2xy, x - y = 6xy**Sol.** x + y = 2xy, x - y = 6xyDividing by xy, we get, $\frac{x}{xy} + \frac{y}{xy} = \frac{2xy}{xy}, \quad \frac{x}{xy} - \frac{y}{xy} = \frac{6xy}{xy}$ $\frac{1}{v} + \frac{1}{x} = 2$...(*i*) $\frac{1}{v} - \frac{1}{x} = 6$...(*ii*) Adding we get, $\frac{2}{v} = 8 \implies 8y = 2$ $\Rightarrow y = \frac{2}{8} = \frac{1}{4}$

Substituting the value of y in (i),

$$\frac{1}{\frac{1}{4}} + \frac{1}{x} = 2 \implies 4 + \frac{1}{x} = 2$$

$$\frac{1}{\frac{1}{4}} + \frac{1}{x} = 2 \implies 4 + \frac{1}{x} = 2$$
Hence $x = a, y = b$ Ans.
$$Q. 28. \quad \frac{3}{x+y} + \frac{2}{x-y} = 3,$$

$$Q. 28. \quad \frac{3}{x+y} + \frac{2}{x-y} = 3,$$

$$\frac{2}{x+y} - \frac{3}{x-y} = \frac{11}{3}$$
Sol.
$$\frac{3}{x+y} + \frac{2}{x-y} = 3,$$

$$\frac{2}{x+y} + \frac{3}{x-y} = \frac{11}{3}$$
Put $x+y=a$ and $x-y=b$, then
$$\frac{3}{a} + \frac{2}{b} = 3 \qquad ...(i)$$

$$\frac{ab^2}{x} + \frac{a^2b}{y} = (a^2 + b^2) \qquad ...(ii)$$
Multiplying (i) by b^2 and (ii) by 1,
$$\frac{b}{a} + \frac{b}{b} = \frac{20}{3}$$

$$\frac{ab^2}{x} - \frac{b^3}{y} = 0$$
$$\frac{ab^2}{x} + \frac{a^2b}{y} = (a^2 + b^2)$$

Subtracting, we get,

 $-\frac{b^3}{v} - \frac{a^2b}{v} = -(a^2 + b^2)$ $-\frac{b}{y}(b^2 + a^2) = -(a^2 + b^2)$ $\Rightarrow \quad -\frac{b}{y} = -\frac{a^2 + b^2}{a^2 + b^2} = -1$ $\frac{b}{-} = 1 \implies y = b$

Substituting the value of y in (i),

$$\frac{a}{x} - \frac{b}{b} = 0 \implies \frac{a}{x} - 1 = 0$$

$$\Rightarrow \quad \frac{a}{x} = 1 \implies x = a$$
Hence $x = a, y = b$ Ans.

$$\frac{1}{4} x = x = x = x$$

$$\Rightarrow \frac{1}{4} x = 2 - 4 = -2$$

$$\Rightarrow \frac{1}{x} = 2 - 4 = -2$$

$$\Rightarrow -2x = 1$$

$$\therefore x = \frac{-1}{2}$$
Hence $x = \frac{-1}{2}, y = \frac{1}{4}$ Ans.
Q. 28. $\frac{3}{x+y} + \frac{2}{x-y} = 3$,
 $\frac{2}{x+y} - \frac{3}{x-y} = \frac{11}{3}$
Sol. $\frac{3}{x+y} + \frac{2}{x-y} = 3$,
 $\frac{2}{x+y} + \frac{3}{x-y} = \frac{11}{3}$
Put $x+y = a$ and $x-y = b$, then
 $\frac{3}{a} + \frac{2}{b} = 3$...(i)
Sol. $\frac{a}{x} - \frac{b}{y} = 0$...(i)
 $\frac{ab^2}{x} + \frac{a^2b}{y} = (a^2 + b^2)$...(ii)
Multiplying (i) by b^2 and (ii) by 1,
 $\frac{5}{a} + \frac{5}{b} = \frac{20}{3}$

95

$$\frac{1}{a} + \frac{1}{b} = \frac{4}{3} \quad \text{(Dividing by 5)} \quad \dots (iii)$$

Subtracting (*ii*) from (*i*),

$$\frac{1}{a} - \frac{1}{b} = -\frac{2}{3}$$
 ...(*iv*)

Adding (iii) and (iv),

$$\frac{2}{a} = \frac{2}{3} \implies \frac{2}{a} = \frac{2}{3}$$
$$2a = 6 \implies a = 3$$

Subtracting (iv) from (iii),

$$\frac{2}{b} = \frac{6}{3} \implies 6 \ b = 6$$
$$b = 1$$

Now substituting the value of a and b,

$$x + y = 3$$
 ...(v)
 $x - y = 1$...(vi)

Adding (v) and (vi),

 $2x = 4 \implies x = 2$ Subtracting, $2y = 2 \implies y = 1$ Hence x = 2, y = 1 Ans.

Q. 29.
$$\frac{22}{x+y} + \frac{15}{x-y} = 5, \frac{55}{x+y} + \frac{40}{x-y} = 13$$

Sol.
$$\frac{22}{x+y} + \frac{15}{x-y} = 5$$
$$\frac{55}{x+y} + \frac{40}{x-y} = 13$$
Put $x + y = a$ and $x - y = b$, then
$$\frac{22}{a} + \frac{15}{b} = 5$$
...(i)
$$\frac{55}{x} + \frac{40}{b} = 13$$
...(ii)
Multiplying (i) by 5 and (ii) by 2,
$$\frac{110}{a} + \frac{75}{b} = 25$$
$$\frac{110}{a} + \frac{80}{b} = 26$$

Subtracting, we get,

D

$$-\frac{5}{b} = -1 \implies \frac{5}{b} = 1$$

0:31.

Substituting the value of b in (i),

$$\frac{22}{a} + \frac{15}{5} = 5 \implies \frac{22}{a} + 3 = 5$$
$$\Rightarrow \frac{22}{a} = 5 - 3 = \frac{2}{1} \implies 2a = 22$$
$$a = \frac{22}{2} = 11$$

Nowing substituting the value of a and b, x + y = 11 ...(*iii*) x - y = 5 ...(*iv*)

Adding we get,

$$2x = 16 \implies x = \frac{16}{2} = 8$$

and subtracting, we get,

$$2y = 6 \implies y = \frac{6}{2} = 3$$

Hence x = 8, y = 3 Ans. Q. 30. If 2x + y = 32 and 3x + 4y = 68, find the value of (x/y).


Q.

96

	$\therefore \frac{x}{y} = \frac{12}{8} = \frac{3}{2}$ Ans.
	·· y 8 2 Alls.
. 31.	If $7x = 10y + 4$ and $12x + 18y = 1$, find
	the values of $(4x + 6y)$ and $(8y + x)$.
Sol.	7 x = 10 y + 4
	$\Rightarrow 7x - 10y = 4 \qquad \dots (i)$
	12 x + 18 y = 1(<i>ii</i>)
•	Multiplying (i) by 9 and (ii) by 5,
	63 x - 90 y = 36
	60 x + 90 y = 5
	Adding, $123 x = 41$
	$x = \frac{41}{123} = \frac{1}{3}$
	$x = \frac{1}{123} - \frac{1}{3}$
	Substituting the value of x in (1),
	$7 \times \frac{1}{3} - 10 y = 4$
	Adding we get. 8
	$\Rightarrow \frac{7}{3} - 10 y = 4$
	3
	$-10 y = 4 - \frac{7}{3} = \frac{5}{3}$
	3 3
	$y = -\frac{5}{3 \times 10} = -\frac{1}{6}$
	Now

Q

32.	The sides of an equilateral triangle are $(x + 3 y)$ cm, $(3 x + 2 y - 2)$ cm and
	$\left(4x+\frac{1}{2}y+1\right)$ cm. Find the length of each side.
Sol.	··· Triangle is an equilateral.
	. Its sides are equal.
	: $x + 3y = 3x + 2y - 2 = 4x + \frac{1}{2}y + 1$
	Taking $x + 3 y = 3 x + 2 y - 2$
	$\Rightarrow 3x + 2y - x - 3y = 2$
	$\Rightarrow 2x - y = 2 \qquad \dots (i)$
	Again taking,
	$3x + 2y - 2 = 4x + \frac{1}{2}y + 1$
	$\Rightarrow 3x + 2y - 4x - \frac{1}{2}y = 1 + 2$
	$\Rightarrow -x + \frac{3}{2}y = 3$
	$\Rightarrow -2x + 3y = 6$
	$\Rightarrow -2x + 3y = 6 \qquad \dots (ii)$
	Adding (i) and (ii),
	2y = 8
	$\Rightarrow \qquad y = \frac{8}{2} = 4$
	Substituting the value of y in (i),
	2x - 4 = 2
	$\Rightarrow 2x = 2 + 4$




 $\left(\frac{1}{6}\right)$

(i) $4x + 6y = 4 \times \frac{1}{3} + 6 \times ($

2x = 6 \Rightarrow $\therefore x = 3$ Now sides of the equilateral triangle are $(3 + 3 \times 4), (3 \times 3 + 2 \times 4 - 2),$ $(4 \times 3 + \frac{1}{2} \times 4 + 1)$ or 15, 15, 15 cm. Ans.

97



```
(ii) x + 5 = 0
x = -5
It is a line parallel to y-axis where x = -5.

(iii) y = 6

It is a line parallel to x-axis where y = 6.

(iv) y + 7 = 0

y = -7

It is a line parallel to x-axis where y = -7.

Downloaded from https:// www.studiestoday.com
```




 $(v) \quad y = x$

By giving different value to x, we find corresponding values of y as given :

x	2	4	-4
у	2	4	-4

Now we will plot the points as shown in graph below : (2, 2), (4, 4) and (-4, -4) on graph and by joining these points, we get the line y = x.



 $(vi) \quad x+y=0 \qquad \implies \qquad x=-y$

By giving different values to y, we get corresponding values of x as given

-6

-7

-8


-9

-10


x	-2	-4	. 3
y	2	4	-3

Now we will plot the points (-2, 2), (-4, 4) and (3, -3) and by joining them, we get the line x + y = 0 as shown in graph above. Downloaded from https:// www.studiestoday.com





Now plot the points (0, 0), (-2, 3) and (-4, 6) and by joining them, we get a line as shown below




100 00

 $x = \frac{9 - 2y}{5}$ (ii) $5x + 2y = 9 \implies 5x = 9 - 2y \implies 3x = 9 - 2y$ Now giving different values to y, we get corresponding values of x. then $x = \frac{9-2\times2}{5} = \frac{9-4}{5} = \frac{5}{5} = 1$ If y = 2, If y = 7, then $x = \frac{9 - 2 \times 7}{5} = \frac{9 - 14}{5} = \frac{-5}{5} = -1$ then $x = \frac{9-2 \times (-3)}{5} = \frac{9+6}{5} = \frac{15}{5} = 3$ If y = -3, 3 1 x -3 ... 7 2 y

Now plot the points (1, 2), (-1, 7) and (3, -3) or the graph and by joining them, we get a line as shown in the graph.

(*iii*) x - 5y + 4 = 0



101

1

By giving different values to y, we get corresponding values of x.

 $x = 5 \times 1 - 4 = 5 - 4 = 1$ If y = 1, then $x = 5 \times 0 - 4 = 0 - 4 = -4$ If y = 0, then $x = 5 \times (2) - 4 = 10 - 4 = 6$ If y = 2, then -4 6 1 x Hence Hence data data 0 2 y

Now plot the points (1, 1), (-4, 0) and (6, 2) and by joining them, we get a line as shown in the graph.

Now plot these points (+ 3, 2); (0, -

(*iv*) 3x - 5y + 1 = 0 \Rightarrow 3 x = 5 y - 1 $x = \frac{5y - 1}{3}$

Now giving different values to y, we get corresponding values of x.

 $x = \frac{5 \times (-1) - 1}{2} = \frac{-5 - 1}{3} = \frac{-6}{3} = -2$ If y = -1, then $x = \frac{5 \times 2 - 1}{3} = \frac{10 - 1}{3} = \frac{9}{3} = 3$ If y = 2, then If y = -4, then $x = \frac{5(-4) - 1}{3} = \frac{-20 - 1}{3} = \frac{-21}{3} = -7$ x -2 3 -7 -1 2 -4 Hence V

Now plot the points (-2, -1), (3, 2) and (-7, -4) on the graph and by joining them, we get a line as shown in the graph.


Q.3. Draw the graph of the equation 4x + 3y + 6 = 0. From the graph, find :

- (i) y_1 , the value of y, when x = 6.
- y_2 , the value of y, when x = -6. (ii)
- 4x + 3y + 6 = 0Sol.

$$4x = -3y - 6$$
$$x = \frac{-(3y + 6)}{4}$$


Now giving some different values to y, we get corresponding values of x.

If
$$y = 2$$
, then $x = \frac{-(3 \times 2 + 6)}{4} = \frac{-12}{4} = -3$
If $y = -2$, then $x = \frac{-[3 \times (-2) + 6]}{4} = \frac{-[-6 + 6]}{$



If
$$y = -6$$
, then $x = -\left[\frac{3 \times (-6) + 6}{4}\right] = -\frac{(-18 + 6)}{4} = \frac{-(-12)}{4} = \frac{12}{4} = 3$
Hence $\frac{x - 3 \quad 0 \quad 3}{y \quad 2 \quad -2 \quad -6}$

Now plot these points (-3, 2), (0, -2) and (3, -6) on the graph and by joining them, we get a line as shown.




From the graph,

- (i) If x = 6, then value of y i.e. y_1 will be = -10.
- (ii) If x = -6, then value of y i.e. y_2 will be = 6.
- Q. 4. Draw the graph of the equation 2x 3y = 5. From the graph, find :
 - (i) x_1 , the value of x, when y = 7.
 - (ii) x_2 , the value of x, when y = -5.


Sol.		$\Rightarrow 2x = 3y + 5$
	$\Rightarrow \qquad x = \frac{3y+5}{2}$ Now giving some different	nt values to y , we get the corresponding values of x .
	If $y = 1$, then	$x = \frac{3 \times 1 + 5}{2} = \frac{3 + 5}{2} = \frac{8}{2} = 4$
	If $y = -1$, then	$x = \frac{3 \times (-1) + 5}{2} = \frac{-3 + 5}{2} = \frac{2}{2} = 1$
	If $y = -3$, then	$x = \frac{3 \times (-3) + 5}{2} = \frac{-9 + 5}{2} = \frac{-4}{2} = -2$
	Hence	$\begin{array}{ c c c c c c c c c c c c c c c c c c c$


Now plot the points (4, 1), (1, -1) and (-2, -3) on the graph and by joining them, we get the line as shown.



104

From the graph (i) If y = 7, then the value of x will be $x_1 = 13$. (*ii*) If y = -5, then the value of x will be $x_2 = -5$. Q. 5. Draw the graph of the equation 3x - y = 4. Find graphically : (i) the value of y, when x = -1. (ii) the value of x, when y = 5. Sol. $3x - y = 4 \implies 3x = y + 4 \implies x = \frac{y + 4}{3}$ Now giving some different values to y, we get corresponding values of x. If y = 2, then $x = \frac{2+y}{3} = \frac{6}{3} = 2$ If y = 5, then $x = \frac{5+4}{3} = \frac{9}{3} = 3$ If y = -1, then $x = \frac{-1+4}{3} = \frac{3}{3} = 1$ 2 3 1 x Hence 5 -1 2 V Now plot the points (2, 2), (3, 5) and (1, -1) on the graph and by joining them, we get the line 10





Downloaded from https://www.studiestoday.com From the graph, When x = -1, then the value of y will be = -7and when y = 5, then the value of x will be = 3. Q. 6. Draw the graph of the line given by the equation 5x + 6y = 30. Use this graph to find the area of the triangle formed by the line and the co-ordinate axes. $x = \frac{30 - 6y}{5}$ \Rightarrow 5 x = 30 - 6 y \Rightarrow **Sol.** 5x + 6y = 30Giving some different values to y, we get corresponding values of x. $x = \frac{30 - 6 \times 0}{5} = \frac{30 - 0}{5} = \frac{30}{5} = 6$ If y = 0, then $x = \frac{30 - 6 \times 5}{5} = \frac{30 - 30}{5} = 0$ If y = 5, then $x = \frac{30 - 6 \times 10}{5} = \frac{30 - 60}{5} = \frac{-30}{5} = -6$ If y = 10, then -6 0 6 x Hence 5 10 0 V Now plot the points (6, 0), (0, 5) and (-6, 10) on the graph and by joining them, we get the line as shown. ords more (-6, 10) 10 9 8.

7.

6

5

4

3

2

B(0, 5)

(1) Let the relation between x and y

106

This line intersects x-axis at 6 at A (6, 0) and y-axis at 5 i.e. at B (0, 5).

 $\triangle AOB$ is formed by the line and the axes.

 \therefore Area of right $\triangle AOB$

$$= \frac{1}{2} \times OA \times OB$$
$$= \frac{1}{2} \times 6 \times 5$$

= 15 square units Ans.

Q. 7. Draw the graph from the table given below :

x	0	4	р	-4
У	3	-3	6	q

- From the graph, find the values of p and q. (*i*)
- State the linear relation between x and y. (ii)
- Sol. We plot the points (0, 3) and (4, -3) and by joining them, we get the required line. From the graph, we see that
 - (i) When y = 6,

then value of x *i.e.* p = -2and when x = -4, then value of y i.e. q = 9

(ii) Let the relation between x and y be


y = ax + b

From the table,

Substituting x = 0, and y = 3, we get $3 = a \times 0 + b$ b = 3

Again substituting


$$x = 4, \quad y = -1$$
$$-3 = 4a + 3$$
$$4a = -6$$
$$a = \frac{-6}{4}$$



 $\Rightarrow 4y = -6x + 12$ $\Rightarrow 6x + 4y = 12$ $\Rightarrow 3x + 2y = 6$ Ans. Q. 8. The graph of a linear equation in x and y passes through the points A (4, 0) and B (0, 3). Draw the graph. Find the value of k, if the graph passes through $\left(k, \frac{3}{2}\right)$. Sol. Plot the points (4, 0) and (0, 3) on the graph and join them to form a line as shown. If $y = \frac{3}{2}$, then x = 2.

 $\therefore k=2$ Ans.





EXERCISE 7 (E)

Solve graphically, the simultaneous equations given below :

Q. 1.
$$x + y = 3$$
, $2x + 5y = 12$
Sol. $x + y = 3$

$$x = 3 - y$$

If $y = 0$, then $x = 3 - 0 = 3$
If $y = 1$, then $x = 3 - 1 = 2$
and if $y = 2$, then $x = 3 - 2 = 1$

100	x	3	2	61
and the second sec	У	0	1	2

Hence

Taking scale 2 cm = 1 unit on both the axes, we plot the points (3, 0), (2, 1) and (1, 2) on the graph. Join them to form a line as shown.

4y = -6x + 12

6x + 4y = 12


Q. 8. The graph of a linear equation in y

the graph Find the value of k, if

If $y = \frac{2}{2}$, then x = 2.

3x+2v=6 Ans


Again 2x + 5y = 122x = 12 - 5y




If
$$y = 2$$
, then $x = \frac{12}{2} = \frac{1}{2} = \frac{1}{2} = \frac{1}{2} = \frac{1}{2}$
If $y = 4$, then $x = \frac{12 - 5 \times 4}{2} = \frac{12 - 20}{2} = \frac{-8}{2} = -4$
Hence $\frac{x + 6}{y + 0} = \frac{1}{2} = \frac{1}$

We plot the points (6, 0), (1, 2) and (-4, 4) and join them to form an other line. We see that these two lines intersect each other at (1, 2). \therefore Solution is x = 1, y = 2 Ans.

Q. 2. x + y = 4, 3x - 2y = 7Sol. x + y = 4 \Rightarrow x = 4 - y




Now we plot the points (4, 0), (2, 2) and (0, 4) on the graph and join them to form a line.



111

Now we plot the points (3, 1), (1, -1) and (-1, -5) on the graph and join them to form a line. We see that these two lines intersect each other at (3, 1). Hence, solution is x = 3, y = 1. Ans. **Q.3.** 2x - 3y = 7, x + y = 1**Sol.** 2x - 3y = 72x = 3y + 7 $x = \frac{3y + 7}{2}$ If y = 1, then $x = \frac{3 \times 1 + 7}{2} = \frac{3 + 7}{2} = \frac{10}{2} = 5$ If y = -1, then $x = \frac{3 \times (-1) + 7}{2} = \frac{-3 + 7}{2} = \frac{4}{2} = 2$ Now we plot the points (If y = -3, then $x = \frac{3 \times (-3) + 7}{2} = \frac{-9 + 7}{2} = \frac{-2}{2} = -$ 0.4.3x - 5y + 1 = 0.2x - y + 3 = 02 5 x Hence -3 -1 1 y 10 9 8-7 6 5 4. 28-34=7 3-(-1, 2)



112

Now we plot the points (5, 1), (2, -1) and (-1, -3) on the graph and join them to get the line. Again x + y = 1 $\Rightarrow x = 1 - y$ If y = 0, then x = 1 - 0 = 1If y = 1, then x = 1 - 1 = 0If y = 2, then x = 1 - 2 = -1

Hence

x	1	0	-1	
У	0	1	2	

Now we plot the points (1, 0), (0, 1) and (-1, 2) on the graph and join them to get another line. We see that these two lines intersect each other at (2, -1).

Hence

Hence solution is x = 2, y = -1 Ans.

Q. 4.
$$3x - 5y + 1 = 0, 2x - y + 3 = 0$$

Sol. $3x - 5y + 1 = 0$

$$\Rightarrow$$


$$x = \frac{5y-1}{3}$$


3x = 5y - 1

If y = -1, then $x = \frac{5(-1) - 1}{3} = \frac{-5 - 1}{3}$

$$=\frac{-6}{3}=-2$$


If y = -4, then $x = \frac{5(-4) - 1}{3} = \frac{-20 - 1}{3} = \frac{-21}{3} = -7$ $5 \times 2 - 1 = 10 - 1 = 9$





Now we plot the points (-2, -1), (-7, -4) and (3, 2) on the graph and join them to form a line. Again 2x - y + 3 = 0 $\Rightarrow 2x = y - 3$ $x = \frac{y - 3}{2}$ If y = 1, then $x = \frac{1 - 3}{2} = \frac{-2}{2} = -1$





Hence


Now plot the points (-1, 2), (0, 3) and (1, 5) and join them to form another line. We see that these two lines intersect each other at (-2, 1). Hence solution is x = -2, y = 1. Ans. **Q.5.** 2x - 3y = 6, 2x + y + 10 = 0Sol. $2x - 3y = 6 \implies 2x = 3y + 6$ If y = -2, then x = -(-2+10) = -8If y = 0, then $x = \frac{3 \times 0 + 6}{2} = \frac{6}{2} = 3$

y

Downloaded from https:// www.studiestoday.com

3

5



Hence

ł				No.
	У	0	-2	-4
	1	and the second		

Now we plot the points (3, 0), (0, -2) and (-3, -4) on the graph and join them to form a line. Again 2x + y + 10 = 0 2x = -(y + 10) $x = -\frac{(y + 10)}{2}$ If y = 0, then $x = \frac{-(0 + 10)}{2} = \frac{-10}{2} = -5$ If y = -2, then $x = \frac{-(-2 + 10)}{2} = \frac{-8}{2} = -4$ If y = -4, then $x = \frac{-(-4 + 10)}{2} = \frac{-6}{2} = -3$

115


x	-5	-4	-3
y	0	-2	-4

Hence

Now we plot the points (-5, 0), (-4, -2) and (-3, -4) on the graph and join them to form another line. We see that these two lines intersect each other at (-3, -4).

 \therefore Solution is x = -3, y = -4 Ans.

Q.6.
$$2x - y + 4 = 0, 3x + 2y - 1 = 0$$



$$\frac{-7}{-8} \frac{1}{\sqrt{7}}$$
If $y = 0$, then $x = \frac{0-4}{2} = \frac{-4}{2} = -2$
If $y = 2$, then $x = \frac{2-4}{2} = \frac{-2}{2} = -1$
If $y = 4$, then $x = \frac{4-4}{2} = 0$
Hence $\frac{x - 2}{y - 0} = \frac{1}{2}$

116

Now we plot the points (-2, 0), (-1, 2) and (0, 4) on the graph and join them to form a line. Again 3x + 2y - 1 = 03x = 1 - 2y

Now we plot the points (-5, 0), (-4, -2) and (-3, -4) or $\frac{\sqrt{2-1}}{2} = x$ and join them to form another line. We see that these two lines intersect each other at $\frac{1}{6-3} = x$.

0.6. 12x+0+

If
$$y = 2$$
, then $x = \frac{1-2(2)}{3} = \frac{1-4}{3} = \frac{-3}{3} = -1$

If
$$y = -1$$
, then $x = \frac{1-2(-1)}{3} = \frac{1+2}{3} = \frac{3}{3} = 1$

If
$$y = -4$$
, then $x = \frac{1-2(-4)}{3} = \frac{1+8}{3} = \frac{9}{3} = 3$

x

y

-1

2

1

-1

Hence

Now we plot the points (-1, 2), (1, -1) and (3, -4) on the graph and join them to form another line.

3

-4

We see that the two lines intersect each other at (-1, 2).

Hence solution is x = -1, y = 2 Ans.

Q. 7.
$$x + 3y = 8$$
, $3x = 2 + 2y$
Sol. $x + 3y = 8$

$$x = 8 - 3 y$$
If $y = 2$, then $x = 8 - 3 \times 2 = 8 - 6 = 2$
If $y = 3$, then $x = 8 - 3 \times 3 = 8 - 9 = -1$
If $y = 4$, then $x = 8 - 3 \times 4 = 8 - 12 = -4$


x	2	-1	-4
y	2	3	4

Hence

Now we plot the points (2, 2), (-1, 3) and (-4, 4) on the graph and join them to form a line. Again 3x = 2 + 2y

$$x = \frac{2+2y}{3}$$

If
$$y = 2$$
, then $x = \frac{2+2\times 2}{3} = \frac{2+4}{3} = \frac{6}{3} = 2$
If $y = 5$, then $x = \frac{2+2\times 5}{3} = \frac{2+10}{3} = \frac{12}{3} = 4$
If $y = -1$, then $x = \frac{2+2(-1)}{3} = \frac{2-2}{3} = 0$



Now we plot the points (2, 2), (4, 5) and (0, -1) on the graph and join them to get another line. We see that these two lines intersect each other at (2, 2).

line. We see that these lines intersect e

Hence solution is x = 1, y = -2 Ans.


2x = -1 - 5y = -(1 + 5y)x = -(1 + 5y)

Q.9. 2x+5y=-1,3x-y=7

	Hence solution is $x = 2$,	y = 2	Ans.		
2. 8.	x + y = -1, y - 2x = -4	4			
Sol.	x + y = -1		in an i		
	x = -1 - y = -(1 +	y)		other a	
	If $y = 1$, then	x = -	(1 + 1)) = -2	
	If $y = 2$, then	x = -	(1+2)) = -3	
	If $y = -1$, then	<i>x</i> = -	(1 - 1)	0 = 0.	
		x	72	-3	0
	Hence	y	1	2	-1

O

Now we plot the points (-2, 1), (-3, 2) and (0, -1) on the graph and join them to form a line.



If x = 3, then $y = 2 \times 3 - 4 = 6 - 4 = 2$


Hence

x	0	2	3
y	-4	0	2

x Q. B. x + y = - 1. y - 2 x = - 4

Now we plot the points (0, -4), (2, 0) and (3, 2) on the graph and join them to form another line. We see that these lines intersect each other at (1, -2). Hence solution is x = 1, y = -2 Ans. Q. 9. 2x + 5y = -1, 3x - y = 7Sol. 2x + 5y = -12x = -1 - 5y = -(1 + 5y) $x = \frac{-(1 + 5y)}{2}$ If y = 1, then $x = \frac{-(1 + 5 \times 1)}{2} = \frac{-(1 + 5)}{2} = \frac{-6}{2} = -3$


119



If
$$y = -3$$
, then $x = \frac{-[1+5\times(-3)]}{2} = \frac{-(1-15)}{2} = \frac{-(-14)}{2} = \frac{14}{2} = 7$

X dament

in the second se	x -3 2 7	
Hence	y 1 -1 -3	
Now we plot the points $(-3, 1)$	(2, -1) and $(7, -3)$ on the graded state $(7, -3)$ on the grad	aph and join them to form a line.
Again $3x - y = 7$		
3x = y + 7		
$x = \frac{y+7}{3}$		
If $y = 2$, then $x = \frac{2}{x}$	$\frac{+7}{3} = \frac{9}{3} = 3$	
If $y = -1$, then $x = -\frac{1}{2}$	$\frac{1+7}{3} = \frac{6}{3} = 2$	Hence



If
$$y = -4$$
, then $x = \frac{-4+7}{3} = \frac{3}{3} = 1$
Hence
 $x = \frac{x}{3} = \frac{3}{2} = 1$
 $y = \frac{1}{2}$

Now we plot the points (3, 2)(2, -1) and (1, -4) on the graph and join them to form another line.

We see that the lines intersect each other at the point (2, -1).

Hence the solution is x = 2, y = -1 Ans.


Q. 10.
$$x + 6y = 11, 2x - 3y = 7$$

Sol.
$$x + 6y = 11 \implies x = 11 - 6y$$

If $y = 1$, then $x = 11 - 6 \times 1 = 11 - 6 = 5$
If $y = 2$, then $x = 11 - 6 \times 2 = 11 - 12 = -1$
If $y = 3$, then $x = 11 - 6 \times 3 = 11 - 18 = -7$

(-7, 3)

x+6y=11



8

7

6

5

4

3.

1.

(5, 1)

(-1, 2)

x	5	-1	-7
у	1	2	3

Hence

121

Now we plot the points (5, 1) (-1, 2) and (-7, 3) on the graph and join them to form a line. Again 2x - 3y = 7 2x = 3y + 7 $x = \frac{3y + 7}{2}$ If y = 1, then $x = \frac{3 \times 1 + 7}{2} = \frac{3 + 7}{2} = \frac{10}{2} = 5$ If y = -1, then $x = \frac{3 \times (-1) + 7}{2} = \frac{-3 + 7}{2} = \frac{4}{2} = 2$ If y = -3, then $x = \frac{3(-3) + 7}{2} = \frac{-9 + 7}{2} = \frac{-2}{2} = -1$ Hence $\frac{x + 5}{y} = \frac{2}{1} = -1$

Now we plot the points (5, 1), (2, -1) and (-1, -3) on the graph and join them to form another line.

If y = 0, then $x = \frac{2 \times 0 - 3}{3} = \frac{-3}{3} = -1$

If y = 3, then $x = \frac{2 \times 3 - 3}{3} = \frac{6 - 3}{3} = \frac{3}{3}$

We see that, these lines intersect each other at (5, 1)

Hence the solution is x = 5, y = 1 Ans.

Q. 11.
$$4x-5y=-4$$
, $3x=2y-3$
Sol. $4x-5y=-4$

$$4x = 5y - 4$$

$$x = \frac{5y-4}{4}$$

If y = 0, then $x = \frac{5 \times 0 - 4}{4} = \frac{-4}{4} = -1$


If
$$y = 4$$
, then $x = \frac{5 \times 4 - 4}{4} = \frac{20 - 4}{4} = \frac{16}{4} = 4$

If
$$y = -4$$
, then $x = \frac{5 \times (-4) - 4}{4} = \frac{-20 - 4}{4} = \frac{-24}{4} = -6$

Hence

x	-1	4	-6
у	0	4	-4

Now we plot the points (-1, 0), (4, 4) and (-6, -4) on the graph and join them to form a line. Again 3x = 2y - 3 $x = \frac{2y - 3}{3}$




Hence

		1	and the	
x	-1	1	3	
у	0	3	6	

Now we plot the points (-1, 0), (1, 3) and (3, 6) on the graph and join them to form another line.

We see that the lines intersect each other at the point (-1, 0)Hence solution is x = -1, y = 0 Ans. **Q. 12.** 5x - 8y = -4, 2x - 7y = 6Sol. $5x - 8y = -4 \implies 5x = 8y - 4$ $x = \frac{8y - 4}{5}$ then $x = \frac{8 \times (-2) - 4}{5} = \frac{-16 - 4}{5} = \frac{-20}{5} = -4$ If y = -2, Downloaded from https:// www.studiestoday.com



If
$$y = 8$$
, then $x = \frac{8 \times 8 - 4}{5} = \frac{64 - 4}{5} = \frac{60}{5} = 12$

y

-2

Hence

Now we plot the points (-4, -2), (5, 3) and (12, 8) and join them to form a line.

3

8

Again
$$2x-7y=6 \implies 2x=6+7y \implies x=\frac{6+7y}{2}$$

If
$$y = 0$$
, then $x = \frac{6+7\times0}{2} = \frac{6}{2} = 3$
If $y = -2$, then $x = \frac{6+7\times(-2)}{2} = \frac{6-14}{2} = \frac{-8}{2} = -4$




If
$$y = -4$$
, then $x = \frac{6+7 \times (-4)}{2} = \frac{6-28}{2} = \frac{-22}{2} = -11$
Hence $\frac{x + 3 + 4 + -11}{y + 0 + -2} = \frac{-22}{2} = -11$

Now we plot the points (3, 0), (-4, -2) and (-11, -4) and join them to get another line. We see that these lines interesect each other at (-4, -2)

Hence solution is x = -4, y = -2 Ans.

Q. 13. Solve graphically, the system of equations x + 2y - 7 = 0, 2x - y + 1 = 0. Find the co-ordinates of the points, where the lines meet y-axis.



Now we plot the points (5, 1), (3, 2) and (1, 3) and join them to form a line.


125


	0=1-0-1-1-0-1-
Again $2x - y + 1 = 0 \implies$	$2x = y - 1 \implies x = \frac{y - 1}{2}$
If $y = 1$, then $x = \frac{1-1}{2} = 0$	y=8-2×3=8-6=2
If $y = 3$, then $x = \frac{3-1}{2} = \frac{2}{2} = \frac{2}{2}$	
If $y = 5$, then $x = \frac{5-1}{2} = \frac{4}{2} = \frac{4}{2}$	2
	x 0 1 2
Hence	v 1 3 5
Now we plot the points $(0, 1)$ $(1, 1)$	3) and $(2, 5)$ and join them to for

Now we plot the points (0, 1), (1, 3) and (2, 5) and join them to form another line. We see that line x + 2y - 7 = 0 intersects y-axis at A and the line 2x - y + 1 = 0 intersects yaxis at B and intersect each other at (1, 3)

: Co-ordinates of A are (0, 3.5) and of B are (0, 1) and solution is x = 1, y = 3 Ans.

- Q. 14. Solve graphically, the system of equations 2x + y = 8, x + 1 = 2y. Find the co-ordinates of the points, where the lines meet x-axis.
 - **Sol.** 2x + y = 8





Again 2x - M + L = 1

We see that line x + 2 v -

Co-ordinates of A arc (0, 3.5

If x = 1, then $y = 8 - 2 \times 1 = 8 - 2 = 6$ If x = 2, then $y = 8 - 2 \times 2 = 8 - 4 = 4$ If x = 3, then $y = 8 - 2 \times 3 = 8 - 6 = 2$

Hence

x	1	2	3
y	6	4	2

Now plot the points (1, 6), (2, 4) and (3, 2) on the graph. Join them to form a line. Again x + 1 = 2y

x = 2y - 1

If y = 1, then $x = 2 \times 1 - 1 = 2 - 1 = 1$ If y = 2, then $x = 2 \times 2 - 1 = 4 - 1 = 3$

If y = 3, then $x = 3 \times 2 - 1 = 6 - 1 = 5$

Hence

x	1	3	5
У	1	2	3

Now plot the points (1, 1), (3, 2) and (5, 3) on the graph and join them to form another line. We see that these lines intersect each other at (3, 2)


 \therefore Solution is x = 3, y = 2.

The line 2x + y = 8 intersects the x-axis at (4, 0) and the line x + 1 = 2, y at (-1, 0) Ans.


Q. 15. Draw the graphs of 2x - y - 1 = 0 and 2x + y = 9 on the same axis. Use 2 cm = 1 unit on both the axes. Write down the co-ordinates of the point of intersection of the two lines.

Sol.
$$2x - y - 1 = 0$$

 $2x = y + 1$
 $x = \frac{y + 1}{2}$
If $y = 1$ then $x = \frac{1 + 1}{2} = \frac{2}{2} = \frac{1 + 1}{2}$



Now plot the points (1, 1), (2, 3) and (0, -1) on the graph and join them to form a line. Again 2x + y = 9 y = 9 - 2xIf x = 1, then $y = 9 - 2 \times 1 = 9 - 2 = 7$ If x = 2, then $y = 9 - 2 \times 2 = 9 - 4 = 5$



Now plot the points (1, 7), (2, 5) and (3, 3) on the graph and join them to form another line.

We see that these lines intersect each other at $\left(\frac{5}{2}, 4\right)$.


Q. 16. Use graph paper for this question. Take 2 cm = 1 unit on both the axes. Draw the graphs of x + y + 3 = 0 and 3x - 2y + 4 = 0 on the same axes. (*i*) Write down the co-ordinates of the point of intersection of the lines. *(ii)*


Sol. x + y + 3 = 0

x = -(y + 3)If y = 0, then x = -(0+3) = -3If y = -1 then x = -(-1+3) = -(2) = -2If y = -3, then x = -(-3 + 3) = 0

Hence

x	-3	-2	0
y	0	-1	-3





Again 3x - 2y + 4 = 0 3x = 2y - 4 $x = \frac{2y - 4}{3}$

If
$$y = -1$$
, then $x = \frac{2 \times (-1) - 4}{3} = \frac{-2 - 4}{3} = \frac{-6}{3}$
If $y = 2$, then $x = \frac{2 \times 2 - 4}{3} = \frac{4 - 4}{3} = 0$
If $y = 5$, then $x = \frac{2 \times 5 - 4}{3} = \frac{10 - 4}{3} = \frac{6}{3} = 2$
Hence $\frac{x - 2}{y - 1} = \frac{2}{5}$

Now we plot the points (-2, -1), (0, 2) and (2, 5) on the graph and join them to form another line.

= +2 south these thinks 2+ =

(iii) White down the co-ordinates

(ii) We see that these lines intersect each other at (-2, -1)Hence solution is x = -2, y = -1 Ans.

(+++++×+

then

►X

9

129


- Q. 17. Taking 2 cm = 1 units on both axes, draw the graphs of x + y + 2 = 0 and 3x 4y = 15 on the same axes. Write down the co-ordinates of the point of intersection of the lines.
 - **Sol.** x + y + 2 = 0
 - x = -(y + 2)If y = 0, then x = -(0 + 2) = -2If y = 1, then x = -(1 + 2) = -3
 - If y = -2, t

then	x = -(1+2) = -3
then	x = -(-2 + 2) = 0


TT		
	011	00
		CC.
		-

X'

•	x = -	(-2 +	2) - (0
	x	-2	-3	0
	y	0	1	-2



(1,-3)



-3

-5

-8.

(-3, -6) -6

Now we plot the points (-2, 0), (-3, 1) and (0, -2) on the graph and join them to form a line. Again 3x - 4y = 15 3x = 4y + 15 $x = \frac{4y + 15}{3}$ If y = 0, then $x = \frac{4 \times 0 + 15}{3} = \frac{0 + 15}{3} = \frac{15}{3} = 5$

130

If
$$y = -3$$
, then $x = \frac{4 \times (-3) + 15}{3} = \frac{-12 + 15}{3} = \frac{3}{3} = 1$
If $y = -6$ then $x = \frac{4 \times (-6) + 15}{3} = \frac{-24 + 15}{3} = \frac{-9}{3} = -3$
Hence $\frac{x + 5}{y + 0} = \frac{1}{3} = -3$


Now we plot these points (5, 0), (1, -3) and (-3, -6) on the graph and join them to form another line. We see that these lines intersect each other at (1, -3). Ans.

same aves. Write

Q. 18. Solve graphically, the system of equations 3x - 5y + 18 = 0, 3x + 5y = 12. Take 2 cm = 2 units on x-axis and 2 cm = 1 unit on y-axis.

Sol.
$$3x - 5y + 18 = 0 \implies 3x = 5y - 18 \implies x = \frac{5y - 18}{3}$$

If $y = 0$, then $x = \frac{5 \times 0 - 18}{3} = \frac{-18}{3} = -6$
If $y = 3$, then $x = \frac{5 \times 3 - 18}{3} = \frac{15 - 18}{3} = \frac{-3}{3} = -1$
If $y = 6$, then $x = \frac{5 \times 6 - 18}{3} = \frac{30 - 18}{3} = \frac{12}{3} = 4$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 6)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$
 $(-6, 0)$



131

Now we plot the points (-6, 0), (-1, 3) and (4, 6) on the graph taking 2 cm = 2 units and join them to form a line.

Again $3x + 5y = 12 \implies 3x = 12 - 5y \implies x = \frac{12 - 5y}{3}$ If y = 0, then $x = \frac{12 - 5 \times 0}{3} = \frac{12 - 0}{3} = \frac{12}{3} = 4$ If y = 3, then $x = \frac{12 - 5 \times 3}{3} = \frac{12 - 15}{3} = \frac{-3}{3} = -1$ If y = 6, then $x = \frac{12 - 5 \times 6}{3} = \frac{12 - 30}{3} = \frac{-18}{3} = -6$ Hence $\frac{x + 4 - 1 - 6}{y + 0 + 3 + 6}$

Now we plot the points (4, 0), (-1, 3) and (-6, 6) on the graph taking 2 cm = 2 unit and join them to form another line. We see that these two lines interesect each other at (-1, 3). Hence x = -1, y = 3 is its solution. Ans.


O. 20. Using I cm = I unit on both the axis, draw - b, graphs of the line

1.5 2- 1= 13

- Q. 19. Solve graphically, the simultaneous equations 2x = y + 3, 3y = 5 x. Take 2 cm = 1 units on both the axes.
 - **Sol.** 2x = y + 3

$$x = \frac{y+3}{2}$$

If y = 1, then $x = \frac{1+3}{2} = \frac{4}{2} = 2$



5-

132

If $y=3$,	then	$x = \frac{3}{2}$	$\frac{+3}{2} =$	$\frac{6}{2} =$	3		
If $y = -1$,	then	$x = \frac{-1}{2}$	+ 3 2	$=\frac{2}{2}=$	= 1		
			x	2	3	1	(C ×
Hence			y	1	3	-1	5

Now we plot the points (2, 1), (3, 3) and (1, -1) on the graph taking 2 cm = 1 unit and join them to form a line.

Again
$$3y = 5 - x$$

 $y = \frac{5 - x}{3}$
If $x = 2$, then $y = \frac{5 - 2}{3} = \frac{3}{3} = 1$
If $x = -1$, then $y = \frac{5 - (-1)}{3} = \frac{5 + 1}{3} = \frac{6}{3}$
If $x = 5$, then $y = \frac{5 - 5}{3} = \frac{0}{3} = 0$
 $x = 2 - 1 = \frac{5}{3}$

Hence

Now we plot these points (2, 1), (-1, 2) and (5, 0) on the graph and join them to form another line. We see that these two lines intersect each other at (2, 1).

2

0

Q. 19. Solve graphically, the simultaneous

Hence solution is x = 2, y = 1 Ans.

Q. 20. Using 1 cm = 1 unit on both the axis, draw the graphs of the line


y

5x - y = 15, 5x - 6y + 10 = 0.


- From the graph, find :
- the co-ordinates of the point of intersection of the lines. (*i*)
- the axes of the triangle between the lines and the x-axis. (ii)

Sol.
$$5x - y = 15 \implies 5x = y + 15 \implies x = \frac{y + 15}{5}$$

If $y = 0$, then $x = \frac{0 + 15}{5} = \frac{15}{5} = 3$
If $y = 5$, then, $x = \frac{5 + 15}{5} = \frac{20}{5} = 4$
If $y = -5$, then $x = \frac{-5 + 15}{5} = \frac{10}{5} = 2$
Hence $\frac{x + 3 + 4}{y + 0 + 5} = \frac{10}{5} = 2$



Now plot the points (3, 0), (4, 5) and (2, -5) on the graph and join them to form a line. \Rightarrow 5 x = 6 y - 10 5x - 6y + 10 = 0Again $x = \frac{6y - 10}{5}$ then $x = \frac{6 \times 0 - 10}{5} = \frac{0 - 10}{5} = \frac{-10}{5} = -2$ If y=0, then $x = \frac{6 \times 5 - 10}{5} = \frac{30 - 10}{5} = \frac{20}{5} = 4$ If y = 5, If y = -5, then $x = \frac{6 \times (-5) - 10}{5} = \frac{-30 - 10}{5} = \frac{-40}{5} = -8$. -2 4 -8 x Hence 5 -5 0 V




Now we plot the points (-2, 0), (4, 5) and (-8, -5) on the graph and join them to form another line. We see that these two lines intersect each other at (4, 5).

These lines intersect x-axis at B (-2, 0) and C (3, 0),

 \therefore AD \perp BC

 $\therefore \text{ Area of } \Delta ABC = \frac{1}{2} \times BC \times AD = \frac{1}{2} \times 5 \times 5 = \frac{25}{2} = 12.5 \text{ sq. units Ans.}$

Q. 21. Determine graphically the co-ordinates of the vertices of the triangle, the equations of whose sides are given below :





If $x = -2$,	then	<i>y</i> = -	<u>-2 +</u> 2	$\frac{8}{2} = \frac{6}{2}$	= 3
If $x = -4$,	then	<i>y</i> = -	<u>-4 +</u> 2	$\frac{8}{2} = \frac{4}{2}$	= 2
		x	0	-2	-4
Hence		y	4	3	2

Now we plot the points on the graph and join them to form one side of the triangle. Again $5y - x = 14 \implies 5y = 14 + x$


3	$y = \frac{14}{5}$	+x				
If $x = 1$,	then	$y = \frac{1}{2}$	<u>14 + 1</u> 5	$\frac{l}{5} = \frac{15}{5}$	-= 3	
If $x = -4$,	then	$y = \frac{1}{2}$	<u>4 - 4</u> 5	$\frac{4}{5} = \frac{10}{5}$) = 2	
If $x = -9$,	then	$y = \frac{1}{2}$	<u>4 - 9</u> 5	$\frac{9}{5} = \frac{5}{5}$	= 1	
		x	1	-4	-9	
Hence		y	3	2	1	

Now we plot these points (1, 3), (-4, 2) and (-9, 1) and join them to form a second side of the triangle.

Now we plot these poir

If $x = 1$, then $y = 1 + 2 \times 1 = 1 + 2 = 3$	
If $x = 0$, then $y = 1 + 2 \times 0 = 1 + 0 = 1$	
If $x = -1$, then $y = 1 + 2 \times (-1) = 1 - 2 = -2$	1
x 1 0 -1	

Hence



Now we plot these points (1, 3), (0, 1) and (-1, -1) and join them to form the third side of the triangle.

We see that these three lines intersect each other at three points A (-4, 2), B (1, 3) and C (2, 5) Ans.


Q. 22. Determine graphically the co-ordinates of the vertices of the triangle, whose equations are given below :

$$y=0, \quad y=x$$

and 2x + 3y = 10.

Sol. (i) y = 0, It x-axis (ii) y = xHere values of x and y are equal.





Now we plot these points (1, 1), (2, 2) and (3, 3) and join them to form the second line. Again 2x + 3y = 102x = 10 - 3y

$$x = \frac{10 - 3y}{2}$$
If $y = 0$, then $x = \frac{10 - 3 \times 0}{2} = \frac{10 - 0}{2} = \frac{10}{2} = 5$
If $y = 2$, then $x = \frac{10 - 3 \times 2}{2} = \frac{10 - 6}{2} = \frac{4}{2} = 2$
If $y = 4$, then $x = \frac{10 - 3 \times 4}{2} = \frac{10 - 12}{2} = \frac{-2}{2} = -1$
Hence $\frac{x + 5}{y} = \frac{2}{2} = -1$

Now we plot these points (5, 0), (2, 2) and (-1, 4) and join them to form the third line. We see that these three lines intersect each other at A (2, 2), B (0, 0) and C (5, 0) Ans.

C(2, 5) Ans

Q. 22. Determine graphically the

and 2x + 3y = 10.

137

...(ii)

EXERCISE 7 (F)

- Q. 1. The sum of two numbers is 53 and their difference is 25. Find the numbers.
- Sol. Let first number = x

and second number = y

Now according to the condition,

 $x + y = 53 \qquad \dots (i)$

and x - y = 25

1 2 ... 70

Adding, we get, 2x = 78

 $x = \frac{78}{2} = 39$

and subtracting, we get

 $2y = 28 \implies y = \frac{28}{2} = 14$

- .:. First number = 39 and and second number = 14 Ans.
- Q. 2. The sum of two numbers exceeds thrice the smaller by 2. If the difference between them is 19, find the numbers.
 - Sol. Let first number = xand second number = yAccording to the conditions,

$$x + y = 3y + 2 \implies x + y - 3y = 2$$

$$x - 2y = 2 \qquad \dots (i)$$

and
$$x - y = 19 \qquad \dots (ii)$$

from (*ii*) x = 19 + y ...(*iii*)

According to the conditions,

x + y = 51 ...(i) 2x - 3y = 12 ...(ii)

Multiplying (i) by 3 and (ii) by 1, we get 3x + 3y = 153

2x - 3y = 12Adding, 5x = 165

 $x = \frac{165}{5} = 33$

Substituting the value of x in (i)

33 + y = 51

 \Rightarrow y = 51 - 33 = 18

Hence larger number = 33 and smaller number = 18 Ans.

- Q. 4. Find two numbers such that the sum of twice the first and thrice the second is 103 and four times the first exceeds seven times the second by 11.
- Sol. Let first number = x and second number = y According to the conditions, 2x + 3y = 103 ...(i) 4x - 7y = 11 ...(ii)

Multiply (i) by 7 and (ii) by 3, we get, 14 x + 21 y = 72112 x - 21 y = 33

Substituting the value of x in (i) $19 + y - 7y = 2 \implies 19 - y = 2$ $\Rightarrow -y = 2 - 19 \implies -y = -17$ $\therefore y = 17$ Substituting the value of y in (iii) x = 19 + 17 = 36Hence numbers are 36, 17 Ans. Q. 3. The sum of two numbers is 51. If the larger is doubled and the smaller is tripled, the difference is 12. Find the numbers. Sol. Let larger number = x and smaller number = y Adding, 26 x = 754 $x = \frac{754}{26} = 29$ Substituting the value of x in (i) $2 \times 29 + 3 y = 103$ $\Rightarrow 58 + 3 y = 103$ $\Rightarrow 3 y = 103 - 58 = 45$ $\therefore y = \frac{45}{3} = 15$ Hence first number = 29 and second number = 15 Ans. Q. 5. Find two numbers such that the sum of thrice of the first and the second is 142

138

and four times the first exceeds the second by 138.

Sol. Let the first number = x

and second number = y

Now according to the conditions,

3x + y = 142		(<i>i</i>)
4x - y = 138	Adding	(ii)

Adding, we get

7x = 280

$$x = \frac{280}{7} = 40$$

Substituting the value of x in (i)

 $3 \times 40 + y = 142$ $\implies 120 + y = 142$

 \Rightarrow y = 142 - 120 = 22

 \therefore First number = 40

and second number = 22 Ans.

- Q. 6. Of the two numbers, 4 times the smaller one is less than 3 times the larger one by 6. Also, the sum of the numbers is larger than 6 times their difference by 5. Find the numbers.
- Sol. Let the larger number = x and the smaller number = y Now according to the conditions,

 $y = \frac{-180}{4} = 45$

 \therefore Larger number = 62

and smaller number = 45 Ans.

Q. 7. If from twice the greater of the two numbers, 45 is subtracted, the result is the other number. If from twice the smaller number, 21 is subtracted, the result is the greater number. Find the numbers.

Sol.	Let larger number = x
	and smaller number $= y$
	2x - 45 = y
	$\Rightarrow 2x - y = 45 \dots (i)$
	2y - 21 = x
	$\Rightarrow \qquad x-2y=-21 \qquad \dots (ii)$
	Adding, we get,
	3x - 3y = 24
	$\Rightarrow x - y = 8$ (<i>iii</i>)
	(Dividing by 3)
	Subtracting, we get,
	x + y = 66 (<i>iv</i>)
	Again adding (iii) and (iv), we get
	$2x = 74 \implies x = \frac{74}{2} = 37$
	Subtracting (iii) from (iv),

3x - 4y = 6...(i) x + y - 6(x - y) = 5 $\Rightarrow x + y - 6x + 6y = 5$ -5x+7y=5...(*ii*) \Rightarrow Multiply (i) by 7 and (ii) by 4, 21 x - 28 y = 42-20x + 28y = 20Adding, we get, x = 62Substituting the value of x in (i) $3 \times 62 - 4y = 6$ \Rightarrow 186 - 4 y = 6 \Rightarrow -4 y = 6 - 186 = -180 Downloaded from https:// www.studiestoday.com

2 y = 58 ⇒ y = ⁵⁸/₂ = 29 Hence larger number = 37 and smaller number = 29 Ans.
Q.8. If three times the larger of the two numbers is divided by the smaller, then the quotient is 4 and remainder is 5. If 6 times the smaller is divided by the larger, the quotient is 4 and the remainder is 2. Find the numbers.
Sol. Let the larger number = x and the smaller number = y We know that, number = Divisor × quotient + remainder Now, according to the conditions,

Q. 10

Sol

139

3x = 4y + 5
$\Rightarrow 3x - 4y = 5 \qquad \dots (i)$
and $6y = 4x + 2$
$\Rightarrow 4x - 6y = -2$
$\Rightarrow 2x - 3y = -1 \qquad \dots (ii)$
(Dividing by 2)
Multiplying (i) by 3 and (ii) by 4, we get
9x - 12y = 15
8x - 12y = -4
Subtracting we get,
x = 19
Substituting the value of x in (i)
$3 \times 19 - 4y = 5 \implies 57 - 4y = 5$
$\Rightarrow -4y = 5 - 57 \Rightarrow -4y = -52$
$\therefore \qquad y = \frac{-52}{-4} = 13$

. .

Hence larger number = 19 and smaller number = 13 Ans.

- Q. 9. If 2 is added to each of the two given numbers, then their ratio becomes 1 : 2. However, if 4 is subtracted from each of the given numbers, the ratio becomes 5 : 11. Find the numbers.
 - Sol. Let the first number = xand the second number = yNow according to the conditions,

Subtracting we get,	
$-x = -34 \implies x = 34$	-
Substituting the value of x in (i)	1.
$2 \times (34) - y = -2 \implies 68 - y = -2$	2
$\Rightarrow -y = -68 - 2 = -70$	
$\therefore y = 70$	
Hence numbers are 34, 70 Ans.	
. The difference between two number	s is
12 and the difference between the squares is 456. Find the numbers.	heir
I. Let the first number $= x$	
and second number = y	
Now according to the conditions,	
x-y=12	(i)
and $x^2 - y^2 = 456$.(<i>ii</i>)
Dividing (ii) by (i)	
$x^2 - y^2$ 456	
$\frac{x^2 - y^2}{x - y} = \frac{456}{12}$	
(x+y)(x-y) 456	
$\Rightarrow \frac{(x+y)(x-y)}{x-y} = \frac{456}{12}$	
\Rightarrow $x + y = 38$	(iii)
Adding (i) and (iii), we get	
$2x = 50 \qquad \Longrightarrow \qquad x = \frac{50}{2} = 25$	
and subtracting (i) from (iii), we get	
26	

 $\frac{x+2}{y+2} = \frac{1}{2} \implies 2x+4 = y+2$ $\Rightarrow 2x-y=2-4$ $\Rightarrow 2x-y=-2 \qquad \dots(i)$ and $\frac{x-4}{y-4} = \frac{5}{11}$ $\Rightarrow 11x-44 = 5y-20$ $\Rightarrow 11x-5y = 44-20$ $\Rightarrow 11x-5y = 24 \qquad \dots(ii)$ Multiplying (i) by 5 and (ii) by 1, we get 10x-5y = -1011x-5y = 24 2 y = 26 ⇒ y = 26/2 = 13. Hence numbers are 25 and 13 Ans.
Q. 11. Find the fraction which becomes 1/2 when its numerator is increased by 6 and is equal to 1/3 when its denominator is increased by 7.
Sol. Let numerator of the fraction = x and denominator = y then fraction = x/y

140

Now according to the conditions,

 $\frac{x+6}{y} = \frac{1}{2} \implies 2x+12 = y \dots(i)$ $\frac{x}{y+7} = \frac{1}{3} \implies 3x = y+7$ $\implies 3x-y=7 \qquad \dots(ii)$ Substituting the value of y from (i) in (ii) $3x-2x-12 = 7 \implies x-12 = 7$ x = 7+12 = 19Substituting the value of x in (i) $2 \times 19 + 12 = y \implies y = 38 + 12 = 50$ $\therefore \text{ Fraction} = \frac{x}{y} = \frac{19}{50} \text{ Ans.}$

Q. 12. A fraction becomes $\frac{1}{2}$ when 1 is subtracted from its numerator and 1 is added to its denominator. Also, it becomes $\frac{1}{3}$ when 6 is subtracted from its numerator and 1 from the denominator. Find the original fraction. Sol. Let numerator of the fraction = x and denominator = y then fraction = $\frac{x}{y}$:. y = 25

Hence original fraction $=\frac{x}{y}=\frac{14}{25}$ Ans.

Q. 13. The denominator of a fraction is greater than its numerator by 9. If 7 is subtracted from both, its numerator and denominator, the fraction becomes $\frac{2}{3}$. Find the original fraction.

Sol. Let numerator of fraction = xand denominator = y

then fraction $=\frac{x}{v}$

Now according to the conditions,

y = x + 9 $\Rightarrow x - y = -9$...(i) and $\frac{x-7}{v-7} = \frac{2}{5}$ \Rightarrow 5x - 35 = 2y - 14 ...(ii) $\Rightarrow 5x - 2y = -14 + 35 = 21$...(iii) from (*i*), x = y - 9Substituting the value of x in (ii) 5(y-9)-2y=21 \Rightarrow 5y-45-2y=21 3y = 21 + 45 = 66 $y = \frac{66}{2} = 22$ Substituting the value of y in (*iii*) x = 22 - 9 = 13 \therefore Original fraction $=\frac{x}{v}=\frac{13}{22}$ Ans. Q. 14. A number consists of two digits, the difference of whose digits is 3. If 4 times the number is equal to 7 times the number obtained by reversing the digits, find the number. Sol. Let unit's digit = x

Now, according to the conditions,

 $\frac{x-1}{y+1} = \frac{1}{2} \implies 2x-2 = y+1$ $\implies 2x-y=1+2$ $\implies 2x-y=3 \qquad \dots(i)$ and $\frac{x-6}{y-1} = \frac{1}{3} \implies 3x-18 = y-1$ $\implies 3x-y=18-1=17 \qquad \dots(ii)$ Subtracting (i) from (ii) x = 17-3 = 14Substituting the value of x in (i) $2 \times 14 - y = 3 \implies 28 - y = 3$ $\implies -y = 3 - 28 = -25$

and ten's digit = ythen number = x + 10 y

141

By reversing the digits, the unit's digit of new number = yand ten's digit = xthen number = y + 10 xNow according to the conditions,

 $y-x = 3 \qquad \dots(i)$ and 4(x + 10y) = 7(y + 10x) $\Rightarrow 4x + 40y = 7y + 70x$ $\Rightarrow 40y - 7y = 70x - 4x$ $\Rightarrow 33y = 66x$

$$x = \frac{33y}{66} = \frac{1}{2}y \qquad \dots (ii)$$

Substituting the value of x in (i)

$$y - \frac{1}{2}y = 3 \implies \frac{y}{2} = 3$$

Substituting the value of y in (ii)

y = 6

 \Rightarrow

$$x = \frac{1}{2} \times 6 = 3$$

- :. Number = $x + 10 y = 3 + 10 \times 6$ = 3 + 60 = 63 Ans.
- Q. 15. A number consists of two digits, the difference of whose digits is 5. If 8 times the number is equal to 3 times the number obtained by reversing the digits, find the number.

22 x = 77 y= 2x = 7y...(*ii*) ...(*iii*) from (*i*), x = 5 + ySubstituting the value of x in (ii) $2(5+y) = 7y \implies 10+2y = 7y$ \Rightarrow 7 $y - 2 y = 10 \Rightarrow$ 5 y = 10 $y = \frac{10}{5} = 2$ Substituting the value of y in (iii) $2x = 7 \times 2 \implies 2x = 14$ $\Rightarrow x = \frac{14}{2} = 7$: Number = $x + 10 y = 7 + 10 \times 2$ = 7 + 20 = 27 Ans. Q. 16. The result of dividing a number of two digits by a number with digits reversed

- is $1\frac{3}{4}$. If the sum of the digits is 12, find the number.
- Sol. Let the unit's digit of the number = xand ten's digit = ythen the number = x + 10 yBy reversing the digits, the unit's digit of the new number = yand ten's digit = xthen number = y + 10 x

Sol. Let unit's digit of the number = x and ten's digits = y then number = x + 10 yBy reversing the digits the unit's digit of new number = y and ten's digit = x \therefore number = y + 10 xNow according to the conditions, x - y = 5 ...(i) and 8 (x + 10 y) = 3 (y + 10 x) \Rightarrow 8 x + 80 y = 3 y + 30 x \Rightarrow 80 y - 3 y = 30 x - 8 x

Now according to the conditions, x + y = 12 ...(i) and $\frac{x + 10 y}{y + 10 x} = 1\frac{3}{4} = \frac{7}{4}$ $\Rightarrow 4x + 40 y = 7 y + 70 x$ $\Rightarrow 40 y - 7 y = 70 x - 4 x$ $\Rightarrow 33 y = 66 x$ $\Rightarrow x = \frac{33}{66} y = \frac{y}{2}$...(ii) Substituting the value of x in (i) $\frac{y}{2} + y = 12 \Rightarrow \frac{3}{2} y = 12$

142

 $\Rightarrow \qquad y = 12 \times \frac{2}{3} = 8$ $\therefore \qquad x = \frac{y}{2} = \frac{8}{2} = 4$

Hence given number = x + 10 y= $4 + 10 \times 8 = 4 + 80 = 84$ Ans.

Q. 17. When a two-digit number is divided by the sum of its digits, the quotient is 8. On diminishing the ten's digit by three times the unit's digit the remainder obtained is 1. Find the number.

Sol. Let unit's digit of the number = xand ten's digit = y

then number = x + 10 y

Now according to the conditions,

 $\frac{x+10\,y}{x+y} = 8$ \Rightarrow x + 10 y = 8 x + 8 y10 y - 8 y = 8 x - x $\Rightarrow 2y = 7x$ $\Rightarrow 7x - 2y = 0$...(i) from (*ii*), y = 3x + 1Substituting the value of y in (i)7x-2(3x+1)=07x - 6x - 2 = 0x = 2: $y = 3x + 1 = 3 \times 2 + 1 = 6 + 1 = 7$ Hence number = $x + 10 y = 2 + 10 \times 7$ = 2 + 70 = 72 Ans. Q. 18. A number of two digits exceeds four. times the sum of its digits by 6 and the number is increased by 9 on reversing the digits. Find the number. Sol. Let units digit of the number = xand ten's digit = ythen number = x + 10 yOn reversing the digit of the given number, the unit's digit of new number = y

and ten's digit = x $\therefore \text{ Number } = y + 10 x$ Now according to the conditions, x + 10 y - 4 (x + y) = 6 $\therefore x + 10 y - 4 x - 4 y = 6$ $\Rightarrow -3 x + 6 y = 6$ $\Rightarrow x - 2 y = -2 \qquad ...(i)$ (Dividing by 3) and x + 10 y + 9 = (y + 10 x) x + 10 y = -9 + y + 10 x $\Rightarrow x - 10 x + 10 y - y = -9$ $\Rightarrow -9 x + 9 y = -9$ $\Rightarrow x - y = 1 \qquad ...(ii)$

Subtracting (*ii*) from (*i*) $-y = -3 \implies y = 3$ $\therefore x - 3 = 1 \implies x = 1 + 3 = 4$ Hence number $= x + 10 \ y = 4 + 10 \times 3$ = 4 + 30 = 34 Ans.

- Q. 19. The sum of the digits of a two-digit number is 12. If the digits are reversed, the new number is 12 less than twice the original number. Find the original number.
 - Sol. Let unit's digit of the number = x

and ten's digit = y then the number = x + 10 yOn reversing the digits, the unit's digit = y and ten's digit = x; then number = y + 10 xNow according to the conditions; x + y = 12 ...(i) and y + 10 x = 2 (x + 10 y) - 12 $\Rightarrow y + 10 x = 2 x + 20 y - 12$ $\Rightarrow 10 x - 2 x + y - 20 y = -12$ $\Rightarrow 8 x - 19 y = -12$...(ii) from (i), x = 12 - y

Q. 2

S

143

Substituting the value of x in (<i>ii</i>)
8(12 - y) - 19y = -12
$\implies 96 - 8y - 19y = -12$
\Rightarrow 96 - 27 y = -12
$\Rightarrow -27 y = -12 - 96 = -108$
-108
$\therefore \qquad y = \frac{-108}{-27} = 4$
$\therefore x = 12 - y = 12 - 4 = 8$
Hence number = $x + 10 y = 8 + 10 \times 4$
= 8 + 40 = 48 Ans.
20. If 11 pens and 19 pencils together cost
Rs. 502; while 19 pens and 11 pencils
together cost Rs. 758, how much 3 pens
and 6 pencils cost together ?
Sol. Let price of one pen = $Rs. x$
and price of one pencil = $Rs. y$
According to the conditions,
$11 x + 19 y = 502 \qquad \dots (i)$
19 x + 11 y = 758(<i>ii</i>)
Adding we get,
30 x + 30 y = 1260
$\Rightarrow \qquad x + y = 42 \qquad \dots (iii)$
(Dividing by 30)
Subtracting (ii) from (i), we get
-8x + 8y = -256
$x - y = 32 \qquad \dots (iv)$

Q

21.	5 kg sugar and 7 kg rice together cost
	Rs. 258 while 7 kg sugar and 5 kg rice
	together cost Rs. 246. Find the total cost
	of 8 kg sugar and 10 kg rice.
ol.	Let price of 1 kg sugar = Rs. x
	and price of 1 kg rice = Rs. y
	Now according to the conditions,
	5 x + 7 y = 258(<i>i</i>)
	7x + 5y = 246(<i>ii</i>)
	Adding (i) and (ii), we get
	12 x + 12 y = 504
	$x + y = 42 \qquad \dots (iii)$
	(Dividing by 12)
	and subtracting (ii) from (i)
	2x - 2y = -12
	$x - y = -6 \qquad \dots (iv)$
	(Dividing by 2)
	Again adding (iii) and (iv)
	$2x = 36 \implies x = \frac{36}{2} = 18$
	22 30 - 2 2
	Subtracting (iv) from (iii)
	$2y = 48 \implies y = \frac{48}{2} = 24$
	$2y = 40 \qquad \Rightarrow \qquad y = \frac{1}{2} = 24$
	Price of 1 kg sugar = Rs. 18
	and price of 1 kg rice = Rs. 24
	Now cost of 8 kg sugar and 10 kg of
2702	rice
	$= R_{s} (8 \times 18 + 10 \times 24)$

(Dividing by - 8)

Again, adding (iii) and (iv),

$$2x = 74 \implies x = \frac{74}{2} = 37$$

and subtracting

 $2y = 10 \implies y = \frac{10}{2} = 5$ $\therefore \quad \text{Price of one pen} = \text{Rs. 37}$ and price of one pencil = Rs. 5 Now cost of 3 pens and 6 pencils $= \text{Rs. } (3 \times 37 + 6 \times 5)$ = Rs. 111 + Rs. 30 = Rs. 141 Ans. = Rs. $(8 \times 18 + 10 \times 24)$ = Rs. 144 + Rs. 240 = Rs. 384 Ans.

- Q. 22. One year ago, a man was four times as old as his son. After 6 years, his age exceeds twice his son's age by 9 years. Find their present ages.
 - Sol. Let present age of son = x years and age of his father = y years According to the conditions, One year ago. Age of son = (x + 1) and age of father = (y + 1) years $\therefore y - 1 = 4(x - 1) \Rightarrow y - 1 = 4x - 4$

144

 $\Rightarrow 4x - y = 4 - 1$ $\Rightarrow 4x - y = 3$...(i) 6 years after, age of son = (x + 6) years and age of father = (y + 6) years $\therefore y + 6 = 2(x + 6) + 9$ \Rightarrow y + 6 = 2x + 12 + 9 = 2x + 21 $\Rightarrow 2x - y = -21 + 6$ $\Rightarrow 2x - y = -15$...(*ii*) Subtracting (*ii*) from (*i*)

$$2x = 18 \quad \Rightarrow \quad x = \frac{18}{2} = 9$$

Substituting the value of x in (i)

 $4 \times 9 - y = 3 \implies 36 - y = 3$ $\Rightarrow -y = 3 - 36$ $\Rightarrow -y = -33 \Rightarrow y = 33$

Hence present age of father = 33 years and age of son = 9 years Ans.

Q. 23. 5 years ago, A was thrice as old as B and 10 years later, A shall be twice as old as B. What are the present ages of A and B?

Sol. Let present age of A = x years and age of B = y years

(7)Age of B = (y - 6) years and $2\left(\frac{1}{5}\right)y - 3y = -1500$ 10 years later, $\Rightarrow \quad \frac{14}{5}y - 3y = -1500$ Age of A = (x + 10) years Age of B = (y + 10) years and According to the conditions, (x-5) = 3(y-5)x-5=3y-15x - 3y = -15 + 5x - 3y = -10 $5x = 7 \times 7500$...(i) and (x + 10) = 2(y + 10) \Rightarrow x + 10 = 2 y + 20 $\Rightarrow x - 2y = 20 - 10 = 10$...(ii) Subtracting (ii) from (i)

 $-y = -20 \implies y = 20$ Substituting the value of y in (i) $x - 3(20) = -10 \implies x - 60 = -10$ x = -10 + 60 = 50

Hence present age of A = 50 years and age of B = 20 years Ans.

- Q. 24. The monthly incomes of A and B are in the ratio of 7 : 5 and their expenditures are in the ratio of 3 : 2. If each saves Rs. 1500 per month, find their monthly incomes.
 - Sol. Let income of A = Rs. x p.m.and income of B = Rs. y p.m.and saving of each one = Rs. 1500 p.m. \therefore Expenditure of A = Rs. (x - 1500)and expenditure of B = Rs. (y - 1500)Now according to the conditions,

$$\frac{x}{v} = \frac{7}{5} \Longrightarrow 5 \ x = 7 \ y \qquad \dots(i)$$

and $\frac{x-1500}{y-1500} = \frac{3}{2}$ \Rightarrow 2x-3000 = 3y-4500 $\Rightarrow 2 x - 3 y = -4500 + 3000$ \Rightarrow 2 x - 3 y = -1500 ...(*ii*) from (*i*), $x = \frac{7}{5}y$ 5 years ago, Substituting the value of x in (ii) Age of A = (x - 5) years $=\frac{14y-15y}{5}=-1500$ $\frac{-y}{5} = -1500 \implies y = 7500.$ Substituting the value of y in (i), $x = 7 \times \frac{7500}{5} = 10500$ Hence income of A = Rs. 10500and income of B = Rs. 7500 Ans.

145

- Q. 25. A 90% acid solution is mixed with 97% acid solution to obtain 21 litres of a 95% solution. Find the quantity of each of the solutions to get the resultant mixture.
 - Sol. Let 90% solution = x litres and 97% solution = y litres According to the conditions,

x + y = 21 ...(*i*) 90% of x + 97% of y = 95% of 21

$$\Rightarrow \frac{90}{100}x + \frac{97}{100}y = \frac{95}{100} \times 21$$

$$\Rightarrow 90x + 97y = 1995 \dots (ii)$$

from (i), x = 21 - ySubstituting the value of x in (ii)

90(21 - y) + 97y = 1995 \Rightarrow 1890 - 90 y + 97 y = 1995 \Rightarrow 7 y = 1995 - 1890 = 105

$$y = \frac{105}{7} = 15$$

Substituting the value of y in (i), $x + 15 = 21 \implies x = 21 - 15 = 6$ Hence 90% solution = 6 litres 97% solution = 15 litres Ans. and

Q. 26. There are two examination halls A and B. If 12 pupils are sent from A to B, the number of pupils in each halls becomes the same. If 11 pupils are sent from B to Subtracting (ii) from (i)

y = 57x - 57 = 24(from (i))x = 24 + 57 = 81

Hence no. of pupils in A room = 81 and no. of pupils in B room = 57 Ans.

- Q. 27. A and B each have a certain number of marbles A says to B. "If you give 30 to me, I will have twice as many as left with you." B replies, "If you give me 10, I will have thrice as many as left with you." How many marbles does each have ?
 - Sol. Let no. of marbles A have = xand no. of marbles B have = yAccording to the conditions,

x + 30 = 2(y - 30) \Rightarrow x + 30 = 2y - 60 $\Rightarrow \quad x-2y = -60 - 30$ $\Rightarrow \quad x - 2y = -90 \qquad \dots (i)$ and 3(x-10) = (y+10) $\Rightarrow 3x - 30 = y + 10$ 3x - y = 10 + 30 = 40...(ii) Multiplying (i) by 1 and (ii) by 2 x - 2y = -906x - 2y = 80Subtracting we get $-5 x = -170 \implies x = \frac{-170}{-5} = 34$ Substituting the value of x in (i)34 - 2y = -90 $\Rightarrow -2y = -90 - 34 = -124$ $y = \frac{-124}{-2} = 62$ Hence A has marbles = 34 and B has marbles = 62 Ans. Q. 28. The present age of a man is 3 years more than three times the age of his son. Three

A, then the number of pupils in A is double their number in B. Find the number of pupils in each room. Sol. Let number of pupils in A hall = xand number of pupils in B = yNow according to the conditions, x - 12 = v + 12x - y = 12 + 12 = 24 ...(i) x + 11 = 2(y - 11)x + 11 = 2y - 22 $\Rightarrow x - 2y = -22 - 11 = -33$(ii)



years hence, the man's age will be 10 years more than twice the age of his son. Determine their present ages.

Sol. Let man's present age = x years and his son's age = y years 3 years hence,

> age of man = (x + 3) years and age of his son = (y + 3) years According to the conditions,

 $x = 3y + 3 \qquad \dots(i)$ x + 3 = 2(y + 3) + 10 $\Rightarrow x + 3 = 2y + 6 + 10$ $\Rightarrow x - 2y = 16 - 3 = 13 \qquad \dots(ii)$ Substituting the value of x from (i) in (ii) $3y + 3 - 2y = 13 \Rightarrow y = 13 - 3 = 10$ Substituting the value of y in (i) $x = 3 \times 10 + 3 = 30 + 3 = 33$ Hence age of man = 33 years and age of his son = 10 years Ans.

Q. 29. The length of a room exceeds its breadth by 3 metres. If the length is increased by 3 m and breadth is decreased by 2 metres, the area remains the same. Find the length and breadth of the room.

Sol. Let length of room = x metres and breadth of room = y metres

- 2(y+3) 3y = -6 $\Rightarrow 2y+6-3y = -6$ -y = -6-6 = -12 $\therefore y = 12$ and x = y+3 = 12+3 = 15Hence length of the room = 15 m and breadth of the room = 12 m Ans.
- Q. 30. The area of a rectangle gets reduced by 8 m², if its length is reduced by 5 m and breadth increased by 3 m. If we increase the length by 3 m and breadth by 2 m, the area is increased by 74 m². Find the length and breadth of the rectangle.
 - Sol. Let length of rectangle = x m and breadth = y m and area = xy m² In first case,

(x-5) (y+3) = xy-8 $\Rightarrow xy+3x-5y+5 = xy-8$ $\Rightarrow 3x-5y = -8+15 = 7$ $\Rightarrow 3x-5y = 7$

and in second case,

(x+3)(y+2) = xy + 74 \Rightarrow xy + 2x + 3y + 6 = xy + 74 $\Rightarrow 2x + 3y = 74 - 6 = 68$...(*ii*) Multiplying (i) by 3 and (ii) by 5, 9x - 15y = 2110 x + 15 y = 340Adding, we get $19 x = 361 \implies x = \frac{361}{10} = 19$ Substituting the value of x in (i) $3 \times 19 - 5y = 7 \implies 57 - 5y = 7$ -5y = 7 - 57 = -56 $y = \frac{-50}{-5} = 10$ \therefore Length of rectangle = 19 m and breadth of rectangle = 10 m Ans. Q. 31. A motorboat takes 6 hours to cover 100 km downstream and 30 km upstream. If

 $\therefore \text{ Area} = \text{length} \times \text{breadth} = x y \text{ m}^2$ In second case,

length = (x + 3) m and breadth = (y - 2) m \therefore Area = (x + 3) (y - 2) m² According to the conditions, x = y + 3 ...(i) (x + 3) (y - 2) = xyxy - 2x + 3y - 6 = xy-2x + 3y = 62x - 3y = -6 ...(ii) Substituting the value of x from (i) in (ii)

the motorboat goes 75 km downstream and returns back to its starting point in 8 hours, find the speed of the motor boat in still water and the rate of the stream.

Sol. Let the speed of motorboat in still water = x km/h

> speed of stream = y km/hand

 \therefore Speed of upstream = (x - y) km/h

and speed of downstream

= (x + y) km/h

Now according to the conditions,

100	30	= 6
x + y	$\overline{x-y}$	- 0

and

 $\frac{75}{x+y} + \frac{75}{x-y} = 8$

Let x + y = a and x - y = b.

$$\therefore \quad \frac{100}{a} + \frac{30}{b} = 6 \qquad \dots (i)$$

$$\frac{75}{a} + \frac{75}{b} = 8$$
 ...(*ii*)

Multiplying (i) by 5 and (ii) by 2

$$\frac{500}{a} + \frac{150}{b} = 30$$
$$\frac{150}{a} + \frac{150}{b} = 16$$

...(*iii*) $\therefore x + y = 25$...(iv)

x - y = 15

Adding, we get,

$$2x = 40, \implies x = \frac{40}{2} = 20$$

and subtracting,

$$2y = 10 \qquad \Rightarrow \qquad y = \frac{10}{2} = 5$$

Hence speed of motorboat in still water = 20 km/h

and speed of stream = 5 km/h Ans.

- Q. 32. A man sold a chair and a table for Rs. 2178, thereby making a profit of 12% on the chair and 16% on the table. By selling them for Rs. 2154, he gains 16% on the chair and 12% on the table. Find the cost price of each.
 - Sol. Let C.P. of chair = Rs. xand C.P. of table = Rs. yIn first case, gain on chair = 12% and gain on table = 16%:. S.P. of chair = $\frac{x(100 + 12)}{100} = \frac{112}{100}x$ and S.P. of table

$$=\frac{y(100+16)}{100}=\frac{116}{100}y$$

In second case, gain on chair = 16% and gain on table = 12%

Subtracting,
$$\frac{350}{a} = \frac{14}{1} \implies 14 \ a = 350$$

 $a = \frac{350}{14} = 25$
Substituting the value of a in (i)
 $\frac{100}{25} + \frac{30}{b} = 6 \implies 4 + \frac{30}{b} = 6$
 $\Rightarrow \frac{30}{b} = 6 - 4 = \frac{2}{1} \implies 2 \ b = 30$
 $b = \frac{30}{2} = 15$

:. S.P. of chair = $\frac{x(100 + 16)}{100} = \frac{116}{100}x$ and S.P. of table $=\frac{y(100+12)}{100}=\frac{112}{100}y$ According to the conditions, $\frac{112}{100}x + \frac{116}{100}y = 2178$ \Rightarrow 112 x + 116 y = 217800 and $\frac{116}{100}x + \frac{112}{100}y = 2154$





...(*ii*) $\Rightarrow 116x + 112y = 215400$ Adding (i) and (ii), we get, 228x + 228y = 433200...(*iii*) x + y = 1900and subtracting (i) and (ii), we get, -4x + 4y = 2400 $\Rightarrow x - y = -600$...(*iv*) Again adding (iii) and (iv), $2x = 1300 \implies x = \frac{1300}{2} = 650$ and subtracting, $2y = 2500 \implies y = \frac{2500}{2} = 1250$ Hence C.P. of chair = Rs. 650 and C.P. of table = Rs. 1250 Ans.

- Q. 33. A man travels 600 km partly by train and partly by car. If he covers 120 km by train and the rest by car, it takes him 8 hours. But, if he travels 200 km by train and the rest by car, he takes 20 minutes longer. Find the speed of the car and that of the train.
 - Sol. Let the speed of car = x km/h

and speed of train = y km/h

Total journey = 600 km.

In first case,

Distance covered by train = 120 km and distance covered by car = 600 - 120= 480 km Time taken = 8 hours In second case, Distance covered by train = 200 km and distance covered by car = 600 - 200= 400 km Total time taken = 8 hours 20 minutes = $8\frac{20}{60} = 8\frac{1}{3} = \frac{25}{3}$ hours. Now according to the conditions

$$\frac{120}{y} + \frac{480}{x} = 8$$

$$\Rightarrow \quad \frac{15}{y} + \frac{60}{x} = 1 \text{ (Dividing by 8)}$$

...(i)

and
$$\frac{200}{y} + \frac{400}{x} = \frac{25}{3}$$

 $\Rightarrow \frac{8}{y} + \frac{16}{x} = \frac{1}{3}$ (Dividing by 25)
...(*ii*)

Multiplying (i) by 8 and (ii) by 15,

80

$$y = 60$$

y

 \therefore Speed of car = 80 km/h and speed of train = 60 km/h Ans.

- Q. 34. 6 men and 8 boys can finish a piece of work in 14 days while 8 men and 12 boys can do it in 10 days. Find the time taken by one man alone and that by one boy alone to finish the work.
 - Sol. Let a man can do the work in = x days and a boy can do the same work in = y days
 - $\therefore \text{ Man's 1 day's work} = \frac{1}{x}$ and boy's 1 day's work = $\frac{1}{y}$

149

...(*ii*)

According to the conditions,

$$\frac{5}{x} + \frac{8}{y} = \frac{1}{14}$$
 ...(i)

and

Multiplying (i) by 3 and (ii) by 2,

 $\frac{8}{x} + \frac{12}{v} = \frac{1}{10}$

$$\frac{18}{x} + \frac{24}{y} = \frac{3}{14}$$
$$\frac{16}{x} + \frac{24}{y} = \frac{2}{10}$$

Subtracting,

$$\frac{2}{x} = \frac{3}{14} - \frac{2}{10} = \frac{30 - 28}{140} = \frac{2}{140} = \frac{1}{70}$$

$$\therefore x = 140$$

Substituting the value of x in (i)

$$\frac{6}{140} + \frac{8}{y} = \frac{1}{14}$$

$$\Rightarrow \frac{8}{y} = \frac{1}{14} - \frac{6}{140} = \frac{10 - 6}{140}$$

$$\Rightarrow \frac{8}{y} = \frac{4}{140} \Rightarrow \frac{8}{y} = \frac{1}{35}$$

$$\therefore y = 8 \times 35 = 280$$
Hence a man can do the work in
$$= 140 \text{ days}$$
and a boy can do the same work in

x + 2y = 100...(ii) Subtracting (i) from (ii) y = 20Substituting the value of y in (i) $x + 20 = 80 \implies x = 80 - 20 = 60$ Hence no. of 25 - P coins = 60and no. of 50-P coins = 20 Ans. Q. 36. A and B together can do a piece of work in 6 days. If A's one day's work be $1\frac{1}{2}$ times the one day's work of B, find in how many days, each alone can finish the work. Sol. (A + B)'s one days work $=\frac{1}{6}$ Let A can do the work in = x days and B can do the work in = y days \therefore A's 1 days' work = $\frac{1}{x}$ and B's 1 day's work $=\frac{1}{v}$ According to the conditions, $\frac{1}{x} = \frac{3}{2} \times \frac{1}{y} = \frac{3}{2y}$...(i)

 $\frac{1}{x} + \frac{1}{y} = \frac{1}{6}$...(*ii*)

Substituting the value of $\frac{1}{x}$ in (*ii*) $\frac{3}{2y} + \frac{1}{y} = \frac{1}{6} \Rightarrow \frac{5}{2y} = \frac{1}{6}$ $\Rightarrow 2y = 30 \Rightarrow y = \frac{30}{2} = 15$ Substituting the value of y in (i) $\frac{1}{x} = \frac{3}{2 \times 15} = \frac{1}{10}$ $\Rightarrow x = 10$ Hence A can do the work in = 10 days and B can do the work in = 15 days Ans.

= 280 days Ans.

Q. 35. A lady has 25-P and 50-P coins in her purse. If in all she has 80 coins totalling Rs. 25, how many coins of each kind does she have ?

Sol. Let no. of 25 - P coins = xand no. of 50 - P coins = yAccording to the condition,

> x + y = 80and $\frac{x \times 25}{100} + \frac{y \times 50}{100} = 25$ $\Rightarrow \frac{x}{4} + \frac{y}{2} = 25$

> > Downloaded from https:// www.studiestoday.com

...(i)