


### Congruence

### Congruence Statement









 $\triangle ABC \cong \triangle FED$  or  $\triangle BCA \cong \triangle EDF$ .


Corresponding angles  $\angle A \cong \angle F$ Corresponding sides


 $\overline{AB} \cong \overline{FE}$ 


 $\angle B \cong \angle E$  $\overline{BC} \cong \overline{ED}$ 

 $\overline{AC} \cong \overline{FD}$ 


**Conditions for Congruence of Two Triangles** 







side are respectively equal



If we gave several people three sticks: 5cm, 7cm and 9cm long, they would all only be able to make the exact sameTriangle.

Q1

#### Answer:

We have to state the correspondence between the vertices, sides and angles of the following pairs of congruent triangles.

(i)  $\triangle ABC \cong \triangle EFD$ 

Correspondence between vertices:

 $A \leftrightarrow E, B \leftrightarrow F, C \leftrightarrow D$ 

Correspondence between sides:

AB = EF, BC = FD, CA = DE

Correspondence between angles:

 $\angle A = \angle E, \angle B = \angle F, \angle C = \angle D$ 

(ii)  $\triangle CAB \cong \triangle QRP$ 

 ${\bf Correspondence\ between\ vertices:}$ 

 $C \leftrightarrow Q, \; A \leftrightarrow R, \; B \leftrightarrow P$ 

Correspondence between sides:

 $\mathit{CA} = \mathit{QR}, \; \mathit{AB} = \mathit{RP}, \; \mathit{BC} = \mathit{PQ}$ 

Correspondence between angles:

 $\angle C = \angle Q$ ,  $\angle A = \angle R$ ,  $\angle B = \angle P$ 

(iii)  $\triangle XZY \cong \triangle QPR$ 

Correspondence between vertices:

 $X \leftrightarrow Q, \; Z \leftrightarrow P, \; Y \leftrightarrow R$ 

Correspondence between sides:

 $XZ = QP, \ ZY = PR, \ YX = RQ$ 

 ${\bf Correspondence\ between\ angles:}$ 

 $\angle X = \angle Q, \ \angle Z = \angle P, \ \angle Y = \angle R$ 

(iv)  $\triangle$  MPN  $\cong \triangle$  SQR

Correspondence between vertices:

 $M \leftrightarrow S, \ P \leftrightarrow Q, \ N \leftrightarrow R$ 

Correspondence between sides:

 $MP = SQ, \ PN = QR, \ NM = RS$ 

 ${\bf Correspondence\ between\ angles:}$ 

 $\angle M = \angle S$ ,  $\angle P = \angle Q$ ,  $\angle N = \angle R$ 

```
Q2
 Answer:
 (i) \triangle ACB \cong \triangle DEF
 (SAS congruence property)
 (ii) \triangle RPQ \cong \triangle LNM
 (RHS congruence property)
 (iii) \triangle YXZ \cong \triangle TRS
 (SSS congruence property)
 (iv) \triangle DEF \cong \triangle PNM
 (ASA congruence property)
 (v) \triangle ACB \cong \triangle ACD
 (ASA congruence property)
Q3
 Answer:
     PL \perp OA
     PM \perp OB
     PL = PM
 To prove:
 \triangle PLO \cong \triangle PMO
 Proof:
 In \triangle PLO \text{ and } \triangle PMO:
 \angle PLO = \angle PMO (90° each)
 PO = PO
                        (common)
 PL = PM
                         (given)
 By RHS congruence property :
 \triangle PLO \cong \triangle PMO
Q4
 Answer:
 Given:
         AD = BC
       AD \parallel BC
 We have to show that AB = DC.
 Proof:
 AD \parallel BC
 \therefore \angle BCA = \angle DAC (alternate angles)
 In \triangle ABC and \triangle CDA:
  BC = DA
                         (given )
 \angle BCA = \angle DAC (proved above)
                         (common)
  AC = AC
 By SAS c ongruence property:
  \triangle ABC \cong \triangle CDA
  =>AB=CD
                                   (corresponding parts of the congruent triangles)
```

```
Q5
 Answer:
 Given:
 AB = AC, BD = DC
 To prove : \triangle ADB \cong \triangle ADC
 Proof:
 (i) In \triangle ADB and \triangle ADC:
 AB = AC
                  (given)
 BD = DC
                  (given)
 DA = DA (common)
 By SSS congruence property:
 \triangle ADB \cong \triangle ADC
 \angle ADB = \angle ADC (corresponding parts of the congruent triangles) ...(1)
 \angle ADB and \angle ADC are on the straight line.
  \therefore \angle ADB + \angle ADC = 180^{\circ}
 \angle ADB + \angle ADB = 180^{\circ}
 => 2\angle ADB = 180^{\circ}
 => \angle ADB = 90^{\circ}
 From (1):
 \angle ADB = \angle ADC = 90^{\circ}
 (ii)\angle BAD = \angle CAD (corresponding parts of the congruent triangles)
Q6
 Answer:
 Given:
 AD is a bisector of \angle A.
 => \angle DAB = \angle DAC
 AD \perp BC
 => \angle BDA = \angle CDA
                            (90° each)
 To prove:
 \triangle ABC is isosceles.
 Proof:
 In \triangle DAB and \triangle DAC:
  \angle BDA = \angle CDA (90° each)
 DA = DA
                          (common)
 \angle DAB = \angle DAC
                         (from 1)
 By ASA congruence property:
  \triangle DAB \cong \triangle DAC
  =>AB=AC (corresponding parts of the congruent triangles)
 Therefore, \triangle ABC is isosceles.
```

```
Q7
 Answer:
 Given:
       AB = AD
       CB = CD
 To prove:
 \triangle ABC \cong \triangle ADC
 Proof:
 In \triangle ABC and \triangle ADC:
 AB = AD
               (given)
 BC = DC
                 (given)
 AC = AC
              (common)
                                            (by SSS congruence property)
 \therefore \triangle ABC \cong \triangle ADC
Q8
 Answer:
 Given:
          PA \perp AB
         QB \perp AB
         PA = QB
 To prove: \triangle OAP \cong \triangle OBQ
 Find whether OA = OB.
 Proof:
 In \triangle OAP and \triangle OBQ:
  \angle POA = \angle QOB (vertically opposite angles)
 \angle OAP = \angle OBQ
                         (90° each)
 PA = QB
                           (g \text{ iven})
 By\ AAS congruence property :
  \triangle OAP \cong \triangle OBQ
  =>OA=OB (corresponding parts of the congruent triangles)
Q9
Answer:
 Triangles ABC and DCB are right angled at A and D, respectively.
 AC = DB
To prove : \triangle ABC \cong \triangle DCB
 In \triangle ABC and \triangle DCB:
 \angle CAB = \angle BDC (90° each)
 BC = BC (common)
 AC = DB
                          (given)
\mathbf{B}y R. H. S. congruence property:
    \triangle ABC \cong \triangle DCB
```

```
Answer:
Given:
\triangle ABC is an isosceles triangle in which AB = AC.
E and F are midpoints of AC and AB, respectively.
To prove:
 BE = CF
Proof:
 E and F are midpoints of AC and AB, respectively.
 => AF = FB, AE = EC
AB = AC
 =>\frac{1}{2}AB=\frac{1}{2}AC
 =>FB=EC
\angle ABC = \angle ACB
                      (angle opposite to equal sides are equal )
 => \angle FBC = \angle ECB
 Consider \triangle BCF and \triangle CBE:
 BC = BC
                      (common)
 => \angle FBC = \angle ECB
 Consider \triangle BCF and \triangle CBE:
 BC = BC
                      (common)
\angle FBC = \angle ECB
                     (proved above)
FB = EC
                      (proved above)
By SAS congruence property:
\triangle BCF \cong \triangle CBE
BE = CF
                (corresponding parts of the congruent triangles)
Q11
Answer:
Given:
AB = AC
 \triangle ABC is an isosceles triangle.
 AP = AQ
To prove:
 BQ = CP
Proof:
 AB = AC (given)
AP = AQ (given)
AB - AP = AC - AQ
 =>BP=CQ
\angle ABC = \angle ACB (angle opposite to the equal sides are equal)
 => \angle PBC = \angle QCB
In \ \triangle \ PBC \ {\rm and} \ \triangle \ QCB :
 PB = QC
               (proved above)
\angle PBC = \angle QCB (proved above)
BC = BC
               (common)
By SAS congruence property:
\triangle PBC \cong \triangle QCB
BQ = CP
                (corresponding parts of the congruent triangles)
```

Downloaded from www.studiestoday.com


```
Answer:
Given:
 ABC is an isosceles triangle.
 AB = AC
BD = CE
To prove:
 BE = CD
Proof:
                           (As, AB = AC, BD = CE)
 AB + BD = AC + CE
 =>AD=AE
 Consider \triangle ACD and \triangle ABE:
AC = AB (given)
\angle CAD = \angle BAE (common)
AD = AE
                (proved above)
By SAS congruence property:
 \triangle ACD \cong \triangle ABE
 =>CD=BE (corresponding parts of the congruent triangles)
Q13
Answer:
 .Given:
 \triangle ABC is an isosceles triangle.
 AB = AC
 BD = CD
 To prove:
 AD bisects \angle A and \angle D.
 Consider \triangle ABD and \triangle ACD:
 AB = AC (given)
 BD = CD (given)
 AD = AD (common)
 By SSS congruence property:
 \triangle ABD \cong \triangle ACD
 => \angle BAD = \angle CAD (by cpct)
 => \angle BDA = \angle CDA (by cpct)
```

#### Q14

#### Answer:

No, its not necessary. If the corresponding angles of two triangles are equal, then they may or may not

They may have proportional sides as shown in the following figure:




#### Q15

#### Answer:

No, two triangles are not congruent if their two corresponding sides and one angle are equal. They will be congruent only if the said angle is the included angle between the sides

Q16

#### Answer:



Both triangles have equal area due to the the same product of height and base. But they are not congruent.

#### Q17

#### Answer:

- (i) the same length
- (ii) the same measure
- (iii)the same side length
- (iv) the same radius
- (v) the same length and the same breadth
- (vi) equal parts

#### Q18

#### Answer:

(i) False

This is because they can be equal only if they have equal sides.

(ii) True

This is because if squares have equal areas, then their sides must be of equal length.

(iii) False

For example, if a triangle and a square have equal area, they cannot be congruent.

(iv) False

For example, an isosceles triangle and an equilateral triangle having equal area cannot be congruent.

(v) False

They can be congruent if two sides and the included angle of a triangle are equal to the corresponding two sides and the included corresponding angle of another triangle.

(vi) True

This is because of the AAS criterion of congruency.

(vii) False

Their sides are not necessarily equal.

(viii) True

This is because of the AAS criterion of congruency.

(ix) False

This is because two right triangles are congruent if the hypotenuse and one side of the first triangle are respectively equal to the hypotenuse and the corresponding side of the second triangle.

(x) True

### Downloaded from www.studiestoday.com